

YMCA Level 3 Technical Specialist in Supporting Participation in Physical Activity: Long-Term Health Conditions (Certificate) (610/6235/2)

Operational start date: 01/08/2026

Qualification Specification



Blank Page

Y Awards Ltd
Suite 14 Conwy House, Castle Court
Swansea Enterprise Park
SA7 9LA
020 3994 9500

www.ymcaawards.co.uk

YMCA Level 3 Technical Specialist in Supporting Participation in Physical Activity: Long-Term Health Conditions (Certificate) (610/6235/2)

Qualification Specification

Copyright © Y Awards Ltd 2026

The content of this document remains the intellectual property of and is copyright to Y Awards Ltd. No part of these materials may be reproduced, stored, copied, edited, or distributed in any form or by any means including photocopying, publishing, recording or by any other electronic or mechanical methods, without the explicit written permission of Y Awards Ltd. Y Awards Ltd reserves the right to seek legal remedies for any such infringement.

Blank Page

Contents

Introduction	1
Aims: Technical specialist qualifications	1
Progression opportunities	3
Stakeholder engagement	4
Entry requirements, prerequisites, and availability	5
Availability	5
Assessment requirement	6
Reasonable adjustments and special consideration	7
Grading and structure	8
Assessment overview	9
Qualification content	21
The role of physical activity and exercise in the prevention and management of long-term health conditions (H/651/1818)	22
Programming, adapting and delivering exercise for adults with long-term health conditions (J/651/1819)	42
Appendix 1: Information sources	62
Appendix 2: Guidance for participant inclusion criteria and scope of practice	66
Appendix 3: Conditions within scope of practice for this qualification	67
Appendix A: Signs, symptoms, pathophysiology and causes	70
Appendix B: Medical interventions, medication side-effects and lifestyle interventions	89
Appendix C: Exercise guidelines and information sources	112
Guidance for training providers	124

Introduction

Y Awards Ltd, has a long and respected history as the first UK awarding organisation to specialise in health and fitness qualifications, launching in 1998 as CYQ.

We're experts in education, health, and wellbeing with over 20 years of experience developing UK-regulated and globally-recognised qualifications. We work closely with industry experts, employers and training providers to make sure that our products and services deliver life-changing opportunities.

With over half a million qualifications awarded, over 400,000 people have advanced their careers with Y Awards Ltd.

Aims: Technical specialist qualifications

Our technical specialist qualifications are designed to:

- build on the outcomes within an occupational standard and enable an individual to specialise in an occupation having achieved the specialist qualification in addition to their attainment of outcomes set out in the standard.
- provide learners with knowledge, skills, and behaviours relevant to developing additional specialist competence in an occupation.
- provide employers with reliable evidence of learners' attainment against occupational outcomes which form the minimum requirements for practice in a specialised role within the occupational area.
- differentiate learner achievement to support employment, progression, and employer selection decisions (where identified as required).
- form part of an engaging course of learning and act as motivation for learners to specialise within the occupational area.

YMCA Level 3 Technical Specialist in Supporting Participation in Physical Activity: Long-Term Health Conditions (Certificate) (610/6235/2)

By completing this qualification, learners will:

- meet Skills England-approved outcomes for clients with long-term health conditions.
- partially meet the requirements as outlined in the Chartered Institute for the Management of Sport and Physical Activity (CIMSPA) population specialism professional standard: 'Working with People with long-term conditions'.

- meet industry requirements to plan, adapt and deliver exercise and physical activity sessions and programmes (within scope of practice) to meet the needs of adults with long-term health conditions.

Long-term health conditions that are covered by this qualification are outlined within Appendix 3: Conditions within scope of practice for this qualification.

Progression opportunities

This qualification is a population specialism. This means it is designed to enable personal trainers to expand their scope of practice to work with adults with long-term health conditions.

This qualification can also lead to further training at other levels to specialise and further increase their scope of practice. For example:

- **Occupational specialism** (to work in additional job roles):
 - YMCA Level 2 Diploma in Exercise and Fitness: Gym Instructor (610/2784/4).
 - YMCA Level 2 Award in Exercise and Fitness: Group Exercise Instructor (610/2792/3).
- **Population specialisms** (to work with a broader range of clients):
 - YMCA Level 2 Award in Engaging Inactive People in Physical Activity to Create Long-Term Behaviour Change (603/7345/3).
 - YMCA Level 2 Award in Engaging Children Aged 0-5 in Sport and Physical Activity (603/7218/7).
 - YMCA Level 2 Award in Engaging Children and Young People in Sport and Physical Activity (603/7216/3).
 - YMCA Level 3 Award in Supporting Participation in Physical Activity: Perinatal (610/0829/1).
 - YMCA Level 3 Award in Supporting Participation in Physical Activity: Disability and Impairments (610/1559/3).
 - YMCA Level 3 Award in Supporting Participation in Physical Activity: Older Adults (610/1668/8).
- **Environment specialisms** (to work in more settings):
 - YMCA Level 2 Award in Developing Sustainable Physical Activity Programmes Within Community Settings (603/7343/X).
- **Lifestyle specialisms** (to support work with a broader range of needs):
 - YMCA Level 2 Award in Supporting Wellness (610/4039/3).
 - YMCA Level 3 Certificate in Supporting Wellness through Lifestyle Behaviour Change (610/4040/X).
- **Technical specialisms** (to work with specific equipment or perform additional roles within the workplace):
 - YMCA Level 2 Award in Mental Health Awareness and Understanding Approaches to Support Individuals (603/7146/8).
 - YMCA Level 3 Award in Emergency First Aid at Work (603/1902/1).
 - YMCA Level 3 Award in First Aid at Work (603/1903/3).
 - YMCA Level 4 Certificate in Advanced Nutrition for Health, Weight Management and Sports Performance (610/2694/3).

Stakeholder engagement

This qualification is mapped against professional standards and meets set criteria and outcomes outlined by the organisations listed below:

- Chartered Institute for the Management of Sport and Physical Activity (CIMSPA).
- Skills England

Qualification	Standard(s)
YMCA Level 3 Technical Specialist in Supporting Participation in Physical Activity: Long-Term Health Conditions (Certificate) (610/6235/2)	Partially maps to V1.0 of the CIMSPA Working with People with long-term conditions, population specialism professional standard. Fully mapped to meet Skills England-approved outcomes for clients with long-term health conditions: approved by Skills England.

Entry requirements, prerequisites, and availability

This qualification has been designed for learners who:

- are 16+ years old.
- hold an occupation entry qualification or apprenticeship endorsed against the CIMSPA Personal Trainer professional standard.
- are able to communicate effectively with individuals and groups.

To have developed teaching skills and confidence, it is recommended that learners have at least six months' experience planning and delivering exercise sessions before taking this qualification.

Availability

Learners can take this qualification in:

Location	Regulated by
England	Ofqual
Wales	Qualification Wales
Northern Ireland	CCEA Regulation
Other UK regions and outside of the UK	Ofqual

Assessment requirement

Client suitability and learner responsibilities for practical assessments.

Learners must have access to real clients who reflect the client profiles identified in the qualification's practical assessment tasks. These clients must present with appropriate criteria as detailed in the relevant assessment brief.

To complete this qualification, learners will require a suitable client for a bespoke client work project (Assessment task 3.1). Learners are required to complete a consultation and assessment with a real client who has:

- two chronic health conditions
- OR.
- one chronic health condition and is physically inactive (not meeting current physical activity guidelines) or is at increased risk of developing chronic health conditions due to physical inactivity and other lifestyle factors.

Sourcing clients


The responsibility for sourcing suitable clients, rests with the learner. However, the centre may provide support where appropriate.

Safeguarding

Should a learner identify their chosen client as vulnerable, they must ensure they are working within their scope of practice and in accordance with safeguarding best practice.

Reasonable adjustments and special consideration

In making these qualifications available, Y Awards Ltd has made every attempt to make sure that there are no unnecessary barriers to achievement. You can find full details of our reasonable adjustment and special consideration policy on our website.



ymcaawards.co.uk/centres/policies-and-procedures

Grading and structure

This qualification is graded Pass or Refer.

A Pass grade demonstrates that a learner has been assessed as fully competent against all assessment criteria within the qualification.

A Refer indicates that a learner has been assessed as not yet competent against one or more of the assessment criteria of the unit and/or qualification. This is a failing grade, and learners will require reassessment to achieve the qualification.


To achieve a Pass, learners must complete the following **two** mandatory units:

Unit number	Unit title	Level	GLH	TQT
H/651/1818	The role of physical activity and exercise in the prevention and management of long-term health conditions	3	120	155
J/651/1819	Programming, adapting and delivering exercise for adults with long-term health conditions	3	100	160

Guided learning hours (GLH): 220

Total qualification time (TQT): 315

Find out more about GLH and TQT on our website:



ymcaawards.co.uk/qualifications/glh-and-tqt

Assessment overview

This qualification is designed to be assessed in line with the learner journey, with learners demonstrating the knowledge, skill, and behaviours outlined in one stage before proceeding to the next. The table below provides details of the tasks within each assessment stage.

Assessment stage and task	Details	Unit(s) assessed
1.1 Long-term health conditions worksheet and professional discussion	There are two parts to this assessment: • 'long-term health conditions' worksheet. • professional discussion 1. Long-term health conditions worksheet. The assessor will allocate a set of seven long-term health conditions. One from each of the following seven categories: • Hypertension OR hypercholesterolemia OR hyperlipidaemia. • Coronary heart disease (post phase IV rehabilitation) OR peripheral vascular disease OR hypothyroidism OR hyperthyroidism • Asthma OR chronic obstructive pulmonary disease (COPD) • Obesity OR type 1 diabetes OR type 2 diabetes • Non-specific low back pain OR osteoporosis • Depression OR stress OR general anxiety disorder • Rheumatoid arthritis OR osteoarthritis OR joint replacement Learners are required to complete the long-term health conditions worksheets for the seven conditions using class notes and independent research from evidence-based reputable sources.	The role of physical activity and exercise in the prevention and management of long-term health conditions (H/651/1818)

Assessment stage and task	Details	Unit(s) assessed
	The completed worksheet template must be submitted to the assessor to mark in preparation for the professional discussion. The assessor will provide feedback within two weeks. 2. Professional discussion Learners will undertake a 10 minute ($\pm 10\%$) professional discussion with their assessor. The aim of the professional discussion is to authenticate the learner's work and confirm their knowledge and understanding relating to the health conditions they were allocated The professional discussion will consist of one broad open-ended question and up to three additional open-ended questions covering the topics listed below: • The role of physical activity and exercise in the prevention and management of long-term health conditions: ○ The wider determinants of health, their impact on health inequalities, and on an individual's health. ○ How specific long-term health conditions (including medical management) may affect participation in exercise and quality of life (giving reference to the short and long-term effects of exercise on the body systems). ○ The benefits, motivators and ways to promote physical activity to overcome barriers to exercise for individuals with long-term health conditions. Learners may refer to their 'Long-term health conditions' worksheet template during the professional discussion. No other notes are permitted.	

Assessment stage and task	Details	Unit(s) assessed
	Further information on assessment is available in the Learner Assessment Record.	
2.1 Applied case study worksheets, professional discussion, and observed practical task	There are three parts to this assessment: 1. applied case study questions. 2. professional discussion plan, deliver, and evaluate. Learners are provided with the following scenario: ‘You have just started to advertise that you are currently training to deliver exercise for clients with long-term health conditions. A range of individuals (see case study briefs) have contacted you asking for information.’ The assessor will select and allocate one set of three case studies (either Set 1, Set 2, Set 3, or Set 4) for the learner to review. 1. Applied case study questions Learners are required to complete the applied case study question template, to: • Summarise the information that may affect each case study’s overall health status, risk stratification (low, medium, high), and readiness to participate in exercise or reasons for exclusion. • Explain any specific health conditions, contraindications or needs that may exceed scope of practice and the action they would take. Once completed, the applied case study question template must be submitted to the assessor to mark in preparation for the professional discussion. The assessor will provide feedback within two weeks.	The role of physical activity and exercise in the prevention and management of long-term health conditions (H/651/1818) and Programming, adapting and delivering exercise for adults with long-term health conditions (J/651/1819)

Assessment stage and task	Details	Unit(s) assessed
	2. Professional discussion Learners will undertake a 12 minute ($\pm 10\%$) professional discussion with their assessor. The aim of the professional discussion is to authenticate the learner's work and confirm their knowledge and understanding relating to the case studies they were issued. The professional discussion will consist of one broad open-ended question and up to four additional open-ended questions covering the topics listed below: Unit: The role of physical activity and exercise in the prevention and management of long-term health conditions (H/651/1818) • LO3: Understand the risks and contraindications to exercise for adults with long-term health conditions: ○ considerations for managing risks for individuals with comorbidities and multiple morbidities. ○ Different risk stratification models used to assess risk of participation. • LO4: Understand guidelines for best practice when delivering physical activity and exercise for adults with long-term health conditions: ○ the role of exercise referral schemes in supporting adults with long-term health conditions. Learners may refer to their applied case study template during the professional discussion. No other notes are permitted. 3. Plan, deliver, and evaluate. On completion of the professional discussion, the assessor will allocate ONE of the case studies and the learner will be required to plan, deliver, and	

Assessment stage and task	Details	Unit(s) assessed
	evaluate a session to meet their needs. The session plan should be 45 – 60-minute duration with a minimum of 30 minutes observed. Learners must complete the following records: • a session overview, including PAR-Q+ and informed consent. • a risk assessment • a session plan: ○ a suitable warm-up, including: – mobility (joint actions). – pulse raising activities/exercises – range of motion stretching ○ main component, including: – cardiovascular training. – muscular fitness training ○ a suitable cooldown and flexibility component • programme of changes over a three-month period All exercises selected are the learners' choice and should provide a balanced whole body training approach covering all components of fitness. The session must be delivered live and in real time and be observed by a qualified assessor. The use of pre-recorded video assessment is not permitted. The practical delivery (and planning and evaluation) may be observed as a full session or continuous and observed in stages by the assessor (so the	

Assessment stage and task	Details	Unit(s) assessed
	various class components – warm-up, main component, and cooldown and stretching) can be completed at different times and over time. Peers can be used for this assessment. On completion of delivery, the assessor will ask learners to complete the self-evaluation record. They will be given 30 minutes in total to complete this task (10 minutes for each session component if delivery is continuous). If each component of session is delivered in stages, you will be given 10 minutes to complete this after each component. Further information on assessment is available in the Learner Assessment Record.	

Assessment stage and task	Details	Unit(s) assessed
3.1 Bespoke client work project Please note: This assessment must be conducted on a 'real' client (not peers).	There are five parts to this assessment: 1. consultation and assessment (with audio or video evidence). 2. planning and programming 3. implementation and evaluation (with video evidence) 4. showcase presentation (with video evidence) 5. professional discussion 1. Consultation and assessment Learners are required to complete a consultation and assessment with a real client who has: • two chronic health conditions OR. • one chronic health condition and inactive (not meeting current activity guidelines), or at higher risk of developing chronic health conditions and other lifestyle factors The client's needs must be within scope of practice. The consultation and assessment should last between 45 and 60 minutes. This must be recorded (video or audio) and submitted as part of the assessment. The client must give their permission and state that they agree to the consultation being recorded. Learners should capture notes during the consultation/assessment in the supplied consultation and assessment record and then write up in full sentences immediately afterwards, where required.	Programming, adapting and delivering exercise for adults with long-term health conditions (J/651/1819)

Assessment stage and task	Details	Unit(s) assessed
	2. Planning and programming Learners are required to use the information gathered during the consultation to plan a full session plan for the first session with the client and an outline of anticipated changes to the programme over a three-month period. Learners must complete the following records: • a session overview, including PAR-Q+ and informed consent. • a risk assessment • session plan: ○ a suitable warm-up, including: – mobility (joint actions). – pulse raising activities/exercises – range of motion stretching ○ main component, including: – cardiovascular training. – muscular fitness training ○ a suitable cooldown and flexibility component • programme of changes over a three-month period All exercises selected are the learners' choice and should provide a balanced whole body training approach covering all components of fitness. The consultation and programming records must be submitted to the assessor to check in preparation for implementing the programme. The assessor will provide feedback within two weeks.	

Assessment stage and task	Details	Unit(s) assessed
	3. Programme implementation and evaluation Learners are required to implement the programme and deliver five sessions with their client or participants. These sessions should be recorded using the programme implementation log. Learners are required to outline: • session dates. • client feedback • reviews of client progress – what changes and improvements One session must be recorded and submitted as part of the assessment. The client must give their permission and state that they agree to the session being recorded. 4. Showcase presentation The showcase presentation can only take place once the learner has implemented their programme for the required five sessions. On completion of the programme implementation, learners are required to record a presentation to showcase their work. The recorded presentation must be 10 minutes ($\pm 10\%$) and include: 1. review and evaluation of the learning and assessment journey, including: • the consultation and assessment process and details of the client: ○ background (health needs and activity levels). ○ goals/aims/motivational factors/barriers.	

Assessment stage and task	Details	Unit(s) assessed
	• programme planning and implementation, and how they supported the client towards the achievement of their goals: ○ description of the programme (content, frequency, and how its effectiveness was monitored). ○ reviews, feedback, any modifications made to the programme. • strategies used to influence both short and long-term behaviour change. • evaluation (what worked well, what challenges they faced, what they would do differently with future clients). • Examples of how the learner will or did build strong relationships with a range of health professionals (both managers and practitioners) who are working in local primary, secondary, or tertiary care services, and the wider workforce including social care, third sector professionals and carers to ensure a smooth customer journey. • Examples of how the learner will use or has used their health knowledge to communicate in a credible manner with health, social care, and third sector professionals at different levels working within the health and social care system. 2. Plans for future work with clients with long-term health conditions. The showcase presentation recording must be submitted to the assessor to mark in preparation for the professional discussion. The assessor will provide feedback within two weeks. 5. Professional discussion	

Assessment stage and task	Details	Unit(s) assessed
	Learners will undertake a 15 minute ($\pm 10\%$) professional discussion with their assessor. The aim of the professional discussion is to authenticate the learner's work and confirm their knowledge and understanding relating to programming, adapting, and delivering exercise to clients with long-term health conditions. The professional discussion will consist of one broad open-ended question and up to five additional open-ended questions covering the topics listed below: • how to screen and assess adults with long-term health conditions prior to participation in physical activity and exercise. • how to plan and adapt exercise for adults with long-term health conditions. • how to deliver and review exercise for adults with long-term health conditions. The bespoke client work project will be centre assessed by an assessor using the checklist provided by Y Awards Ltd. Further information on assessment is available in the 'Learner Assessment Record'.	

Using this document

The following pages provide the unit content for this qualification. Each unit includes learning outcomes, assessment criteria, and relevant content for delivery. These are set out below.

Learning outcome ('The learner will')	
Assessment criteria (‘The learner can’) What a learner is expected to know, understand, or be able to do following their learning.	Relevant content (Additional delivery guidance) Suggestions on depth and breadth of content to cover. Learners may not be assessed against all the relevant content identified.

Qualification content

The role of physical activity and exercise in the prevention and management of long-term health conditions (H/651/1818)

Unit aim

To provide the knowledge of the role of physical activity in the prevention and management of long-term health conditions.

Learners will know:

- the signs, symptoms, and progression of specific health conditions.
- the causes and prevalence of specific health conditions.
- how health conditions are treated and managed.
- contraindications and risks for exercise.
- the benefits of referral to exercise and stages of the referral process.

Please note: To support the delivery and assessment of LO1, long-term health condition tables are provided in the appendices at the end of this document

- **Appendix A (p. 71):** Signs, symptoms, pathophysiology, and causes.
- **Appendix B (p. 89):** Medical interventions, medication side-effects, and lifestyle interventions.
- **Appendix C (p. 112):** Exercise guidelines and information sources.

Content


1. Understand long-term health conditions and their effect on health, wellbeing and participation in physical activity and exercise	
1.1 Describe the signs, symptoms and disease progression for specified long-term health conditions	• See information resource: Long-term health conditions tables: ‘Appendix A: Signs, symptoms, pathophysiology, and causes’: ○ Signs and symptoms and pathophysiology/disease progression of the following health conditions: – Hypertension – Hypercholesterolaemia – Hyperlipidaemia – Hypothyroidism

	– Hyperthyroidism – Coronary heart disease (post-phase IV rehabilitation) – Peripheral vascular disease – Diabetes type 1 and 2 – Obesity – Chronic obstructive pulmonary disease (COPD) – Asthma – Osteoarthritis – Rheumatoid arthritis – Osteopenia – Osteoporosis – Low back pain – Joint replacement – Depression – Stress – General anxiety disorder NB: See Appendices 2 and 3 for guidance on scope and inclusion. ○ With consideration to changes to the body systems appropriate to specific conditions (progressive and over time): – Skeletal, i.e. bone density, joint mobility, range of motion (ROM), alignment, and posture. – Neuromuscular, i.e. posture, pelvic floor, power, strength, endurance, sarcopenia, heat production, immune function, fine motor control, coordination, balance and falls risk, reaction time (including ability to respond to instructions), movement speed, sight and hearing, short-term memory. – Cardiovascular, i.e. reduced anaerobic threshold, reduced exercise tolerance, and ability to sustain activity. – Respiratory, i.e. breathlessness. – Cognitive, short-term memory. – Psychosocial, i.e. low mood, low energy, loss of interest in life: ○ Consider the following for all conditions: – Age of onset. – Management.
--	--

	– Individual attitude and response to diagnosis, including cognitive processing time (locus of control). – The importance of mental health and wellbeing for people at high(er) risk of developing, or living with, single LTCs or comorbidities, and the factors that can influence an individual's wellbeing.
1.2 Outline the risk factors and causes for specific long-term health conditions	• Genetics/heredity factors. • The effects of ageing. • The importance of mental health and wellbeing for people at high(er) risk of developing, or living with, single LTCs or comorbidities and the factors that can influence an individual's wellbeing. • Other lifestyle related risk factors: ○ Physical inactivity – identified as one of the top ten causes of diseases and disabilities in England and is responsible for one in six deaths in the UK (Sport England, 2022). ○ Smoking. ○ Alcohol misuse. ○ Unhealthy eating. • Modifiable and non-modifiable risk factors.
1.3 Describe the wider determinants of health, their impact on health inequalities and on an individual's health	• Wider health determinants – economic, social, and environmental, and inequalities as outlined by the World Health Organization (WHO). • Wider or social determinants of health are the various non-medical factors that affect health outcomes, such as social, economic, environmental, political, commercial, and cultural factors. • Some examples of wider 'determinants' would include: ○ The number of fast-food outlets on the high street, which has the potential to influence levels of obesity and diabetes. ○ The availability or work opportunities in a specific location. ○ The accessibility or availability of services to support health in a location. ○ Poverty levels.

- Health inequalities are unfair and avoidable differences in health across the population and between different groups within society. These include how long people are likely to live, the health conditions they may experience and the care that is available to them (NHS England).

Determinants of Health (Dahlgren and Whitehead model) from: Public Health England (2017).



Can impact:

- Longevity of life, life expectancy.
- Healthcare accessible and available.
- The prevalence of some health conditions, e.g. some populations are at greater risk of some conditions.

1.4 Describe how specific long-term health conditions are medically treated and managed

- See AC1.1 for the list of specific conditions.
- See information resource: Long-term health conditions tables: Appendix B: Medical interventions, medication side-effects and lifestyle interventions.
- Use of National Institute for Health and Care Excellence (NICE) guidance for specified conditions, including activity recommendations where specified.
- Medical management – surgery, medication with consideration to:
 - Desired effects of medication or other treatment, e.g. surgery.
 - Side effects of medication or other treatment, e.g. surgery
 - Effects on the exercise response, e.g. heart rate, blood pressure, balance, exercise tolerance, functional capability, energy levels, etc
 - Credible sources to gather information about medication (British National Formulary [BNF] and Monthly Index of Medical Specialities [MIMS])
 - Effects of some treatments and medication on the quality of life, including functional and mental capacity

1.5 Explain lifestyle changes recommended to assist the prevention and management of specific long-term health conditions	• See information resource: Long-term health conditions tables: Appendix B: Medical interventions, medication side-effects, and lifestyle interventions. • General/lifestyle recommendations for specific conditions (use NICE, British Heart Foundation (BHF) and other credible and reliable sources): ○ the Eatwell Guide for healthy adults. ○ Current chief medical officer's (CMO) guidance for physical activity ○ Wellbeing and wellness practices, including mindfulness, breathing exercises, stress management, and cognitive behavioural therapy (CBT) ○ Smoking cessation
1.6 Identify the prevalence of specific long-term health conditions	• Percentage of population with specific long-term health conditions (use of current statistics). • With consideration to: ○ Prevalence of conditions for different groups: – Sex. – Age. – Ethnicity. – Disabilities – physical, visual, hearing: ○ The effects of wider health determinants – social, economic, and environmental factors (non-medical). ○ The number of years lived without a chronic health condition has reduced: – Age 62 years for men. – Age 60 years for women: ○ More people have more than one health condition: – One third of all adults in their late 40s have two or more chronic health conditions (Hall, The Guardian, 2021). ○ The ageing population in the UK is increasing. ○ Multiple health conditions increase the risk of frailty in later life, which can lead to loss of independence, reduced quality of life and increased risk of hospital admissions and death. Reference sources listed in appendix: ○ Centre for ageing better. ○ NHS

	○ Hall, The Guardian (2021)
1.7 Explain how specific long-term health conditions may affect participation in exercise	• With consideration to: ○ Specific health conditions. ○ Associated risk factors – see LO3.2. ○ Effect of ageing, injury, and disease on functional capacity and health outcomes. ○ Individual attitude and their locus of control. ○ The physiology and psychology of pain and how this may impact upon the functional and mental capacity of an individual. • Effects on: ○ Suitability of specific types of activity and exercise. ○ Session planning and design – unit 2. ○ Session delivery – units 2 and 3. ○ Risk stratification – see screening and assessment (unit 2) and impact on: – Scope of practice. – Level of instruction and supervision. – Need for specialist support and supervision for those with more declining health, increased age, and/or significant loss of capacity (higher risk). – Higher level of support may include physiotherapy teams/specialist teams, postural stability instructor (PSI). ○ Appropriate exercise settings – clinical, leisure facilities (studio, gym), community, home-based etc.
1.8 Explain the impact of specific long-term health conditions, their medical management and associated risk factors on quality of life	• Functional movement and ability to perform activities of daily living • Mental health • Longevity of life • Quality of years lived

2. Understand the role of physical activity and exercise in the prevention and management of long-term health conditions	
2.1 Identify guidelines and recommendations for physical activity and exercise for specific long-term health conditions	• Use of ACSM guidelines and recommendations. • Use of NICE guidance and recommendations. • Use of health charity specific guidance and recommendations. • See ‘Appendix 1: Information sources’. NB: It is recommended that as part of delivery, centres include examples of practical sessions delivered for (and ideally with) individuals with health conditions. • See information resource: Long-term health conditions tables: Appendix C: Exercise guidelines and information sources
2.2 Explain the benefits of physical activity and exercise to support successful ageing and assist with the prevention and management of specific long-term health conditions	• Effects of ageing, physical inactivity, and long-term health conditions on anatomy, physiology, and biomechanics. • The protective effects of physical activity and exercise for both the prevention and management of specific long-term health conditions: ○ Difference between physical activity and exercise. ○ Reference to the current Chief Medical Officer’s (CMO) guidance. ○ Successful and active ageing and quality of life. • How exercise can support successful ageing and benefit management of specified long-term health conditions (NICE guidance). • With consideration to: ○ The physiological, biomechanical, anatomical, and psychological response to physical activity: – Short-term and immediate effects – increase to heart rate and breathing rate, muscle temperature, blood pressure, etc. – Long-term effects – stronger heart, increased stroke volume, regulate blood pressure, increase bone density, etc. – Effects associated with different types of exercise and components of fitness. ○ Longer-term benefits: – Maintaining independence.

	– Improving fitness and maintaining ability to perform activities of daily living and continue leisure pursuits. – Supporting the management of existing health conditions and minimising effect on functioning. – Maintaining psychological wellbeing. – Maintaining and building social connections.
2.3 Explain the barriers to physical activity and exercise for adults with long-term health conditions and how to overcome these	• Awareness of segmentation tools to support the understanding of barriers, motivators, and activity preferences. • Identification of populations most at risk of physical inactivity and other health inequalities. • Real and perceived barriers: ○ Psychological barriers – lack of confidence, self-efficacy, self-esteem, mental health (depression, anxiety), fear of making health conditions or injuries worse, fear of being too old or not fit enough etc. ○ Physical/environmental barriers – accessibility and availability of appropriate activities, cost, location, method of booking sessions (such as use of apps etc.) transport links, lack of transport, suitability of exercise sessions, suitability of instructors. ○ Barriers/concerns linked with health: – Medical diagnosis and conditions, e.g. osteoarthritis etc. – Medications and their effects. – Comorbidities and multimorbidity. – Functional, strength and balance – falls risk and fear of falling, mobility issues, vision, and hearing. – Cognitive impairments – memory, responding to instructions, etc. • How to support individuals and overcome barriers to promote active ageing. • The importance of mental health and wellbeing for people with a high(er) risk of developing or living with long-term conditions and the factors that can influence an individual's wellbeing.
2.4 Explain motivators for physical activity and exercise for adults with long-term health conditions	• Consideration of behaviour change models. • Awareness of segmentation tools to support the understanding of barriers, motivators, and activity preferences. • Specific motivators:

	○ Recommendation of GP. ○ Support of family and friends. ○ Personal motivators – self-determination, gender, life stage. ○ Positive mental health as a motivator. ○ Maintaining independence and functioning. ○ Reduce risk of chronic health conditions. ○ Management of chronic health conditions. ○ Management of mental health and wellbeing. ○ Improve quality of life. ○ Socialisation and structure for day – getting out of house.
2.5 Outline appropriate methods to promote exercise, physical activity, and wellbeing initiatives to support adults with long-term health conditions	● The importance of providing a full spectrum of opportunity for clients to engage in physical activity that includes activities of daily living, active travel, active recreation through to structured exercise, dance, and sport. ● With consideration to emerging messages regarding ‘reframing the exercise message’ and the role of non-exercise activity thermogenesis (NEAT) – all daily activities that are not exercise, or sport, or sleeping and eating. ● The principles of self-management which enable participants to maintain physical activity and other lifestyle changes beyond the intervention. ● The role of evidence-based technologies that support the uptake and maintenance of physical activity. ● Other health interventions and other wider community assets that can support the participant. ● Knowing how, when, and where to signpost, e.g. health trainers, improving access to psychological therapies (IAPT), smoking cessation, physiotherapy, pharmacy. ● Support services – GP, charities, exercise referral services, leisure services. ● Local physical activity opportunities to signpost individuals onward. ● Range of promotional and marketing strategies. ● Link to motivators – see above – AC 2.4 Explain motivators for physical activity and exercise for adults with long-term health conditions. ● Appropriate images – positive, inclusive. ● Helpful language and messages.

	• Consider targeted and specific marketing for gender, ethnicity, socio-economic status, health status, identity and how individuals identify. • How to communicate the impact of exercise on the human body in simple terms to a participant. • Peer mentoring. • Ongoing and emerging research and initiatives to support lifestyle and behaviour change interventions: ○ One-to-one, group and peer behaviour change support. ○ Face to face versus remote communication, e.g. digital, print and telephone. ○ Combined interventions to meet the needs of individuals with long-term health conditions, including those with health inequalities. ○ The use of customer relationship management (CRM) systems to monitor change in behaviour and to track and tailor follow up motivational support based on individual need(s) and circumstance.
--	--

3. Understand the risks and contraindications to exercise for adults with long-term health conditions

3.1 Identify the contraindications to exercise	• Reference to the current American College of Sports Medicine (ACSM) guidelines for specific conditions (PAR-Q+ and Algorithm). • Use of NICE guidance (condition specific recommendations and guidelines). • Use of information to guide recommendations on participation/deferral/referral and referral sources, e.g. GP. • Absolute contraindications for exercise (general) – use ACSM guidelines for exercise testing and prescription, such as: ○ Any uncontrolled or unstable condition, e.g. not managed by medication. ○ Resting systolic blood pressure at (or above) 180mmHg/DBP 100mmHg (BHFNC, 2010). ○ Uncontrolled resting tachycardia at or above 120 bpm. ○ Experiences a negative change or increase in pain during exertion. ○ Dizziness or excessive breathlessness during exertion. ○ Significant postural hypotension. • Absolute contraindications specific to different conditions. Reference sources (listed at end):
---	--

	○ American College of Sports Medicine (ACSM) Guidelines for Exercise Testing and Prescription. ○ BHFNC toolkit ○ NICE guidelines (condition specific) ○ ACSM Exercise Management for Persons with Chronic Diseases and Disabilities
3.2 Explain the risks associated with participation in exercise for specific long-term health conditions and ways to manage these risks	● Risks (accidents and emergencies) associated with specific long-term health conditions: ○ Hypoglycaemia. ○ Strains and sprains. ○ Exercise induced asthma (EIA). ○ Breathlessness. ○ Joint pain and discomfort. ○ Loss of balance falls risk and fractures. ○ Cardiac incidents – angina, heart attack. ○ Dizziness and fainting. ○ Cramps. ○ Hyperthermia. ○ Dehydration. ○ Myocardial infarction. ● Condition negatively impacted/worsening. ● Reduction in self-esteem. ● Effects of common functional impairments that increase risk of participation: ○ Risk of falling or fear of falling. ○ Hearing, speech, and sight impairments. ○ Memory or cognitive decline. ○ Proprioception. ○ Soft tissue injury. ○ Joint pain. ○ Fatigue. ○ Grip. ○ Range of motion. ○ Motivation. ○ Muscle strength. ○ Reaction time.

	○ Ability to transition from floor to stand. ○ Ability to perform certain exercises. ○ Ability to achieve specific workloads. ○ Energy levels and effort required to complete specific tasks. ● The importance of contingency planning to manage any identified risks, which may include deferral or referral of exercise and signposting to specialist services, e.g. falls prevention service.
3.3 Outline the side effects of common pharmaceutical treatments and their implications for exercise tolerance and functional capability	● Effects of specific medication(s) may include alterations to: ○ Alertness. ○ Posture. ○ Balance. ○ Falls risk. ○ Proprioceptive feedback. ○ Exercise intensity and heart rate and blood pressure response. ○ Urinary urgency and frequency. ○ Postural hypotension. ○ Masked pain levels. ○ Coordination and reaction time.
3.4 Explain how multiple health conditions and medications may affect risk stratification and readiness to participate	● Effects on risk stratification and scope of practice: ○ When multiple health conditions are present, this may increase risk stratification. ○ Individuals with higher or high-risk stratification are outside of scope of practice. ○ Multiple medications (4+) increases risk of falls in older adults.
3.5 Outline considerations for managing risks for individuals with comorbidities and multiple morbidities, including reasons for exclusion	● See LO1 for risks. ● See AC4.2 for exclusion criteria. ● Define comorbidities and multiple morbidities. ● Awareness of absolute contraindications. ● Signposting to other professionals if needs exceed scope. ● Requirement for higher level qualifications, depending on risk stratification. ● With consideration to own ability, competence, and confidence to:

	○ Manage any potential medical emergency/risks associated with conditions. ○ Consider all specific conditions in exercise planning. ○ Adapt exercise to meet individual needs. ○ Apply the principles of training to modify and adapt the exercise (if within scope of practice). Information sources: • Richmond Group of Charities (2018) Multimorbidity Report listed in the references.
3.6 Evaluate different risk stratification models used to assess risk of participation	• PAR-Q+ and e-PARmed-x + • ACSM algorithm • Irwin and Morgan • Health and Safety Executive (HSE) – risk assessment tool (not condition specific) • With consideration to other specific questionnaires, such as: ○ Falls risk and balance (FRAT and CONFbal) ○ Back pain questionnaires (Oswestry and Roland-Morris etc.) ○ Mental Health (PHQ-9 etc.) NB: Centres may evaluate a broader range of additional questionnaires. Those listed are shared as examples.

4. Understand guidelines for best practice when delivering physical activity and exercise for adults with long-term health conditions

4.1 Identify a range of evidence-based information sources to support working with adults with long-term health conditions	• Evidence-based information sources: ○ Chief Medical Officer (CMO) guidance – most current – 2019. ○ American College of Sports Medicine (ACSM). ○ Research studies on specific long-term health conditions. • Other information sources: ○ Public Health England – healthy eating and stop smoking brief, change4life. ○ Association for Nutrition (AfN) – competence framework – healthy eating and nutrition. (see 4.2). • GP and other healthcare services (such as physiotherapists): ○ Local and national charities – specific health conditions, e.g. British Heart Foundation, Diabetes UK, Asthma UK.
---	---

	The benefits of working with other services to support participation in physical activity. • Uses: ○ To support health outcomes. ○ To assist activity planning. ○ To inform own understanding. See 'Appendix 1: Information sources'.
4.2 Explain own working role and responsibilities and the importance of working within scope of practice and the boundaries of personal and occupational competence	• Know how to liaise with health care and social care professionals where there are considerable concerns for a participant's safety and refer where necessary. • Refer to roles and responsibilities outlined in the exercise referral toolkit (2010). • Importance of working in scope: • Client safety and wellbeing. • Maintain professional practice and role boundaries. • Adhere to insurance guidance. • Adhere to safeguarding legislation. • With consideration to industry code of conduct, ethical issues, and other associated codes of conduct (such as Association for Nutrition – AfN) relevant to the role in terms of both relationships with clinicians, nutritionists/dieticians, referring professionals, and participants: ○ Fundamentals of human nutrition ○ Improving health and wellbeing ○ Nutrition monitoring and data collection techniques ○ With consideration to scope of practice as outlined by AfN • Inclusions for this qualification: ○ Conditions listed at 1.1. ○ Stable and controlled only and not in progressed stages, or with complications • Exclusions: requirement for additional qualifications: ○ Conditions not listed at 1.1. ○ Any condition in more progressed stage. ○ Adults with multiple health conditions, higher risk stratification, and declining capacity. ○ Frailer and dependent older adults – falls and fracture risk – significant loss of capacity (Postural stability instructor -PSI).

- Cardiovascular events – stroke, heart attack, angina (British Association of Cardiovascular Prevention and Rehabilitation – BACPR).
- Respiratory conditions – severe and very severe.
- Mental health conditions – more severe and enduring (eating disorders, substance misuse, bipolar, schizophrenia, etc.).
- Cancer.
- Spinal cord injury.
- Connective tissue disease.
- Neurological conditions (multiple sclerosis, Parkinson's disease, dementia, fibromyalgia, chronic fatigue syndrome).
- If in doubt, lacking confidence regarding skills and knowledge (or the appropriateness of the content and structure of a specific type of session), this may also be reason for exclusion.

Please note:

- Risk stratification for participation in exercise is currently determined using different tools, e.g. ACSM algorithm, PAR-Q+ and Irwin and Morgan:
 - The level of risk in an exercise and activity environment will be impacted by several factors, including:
 - The risk stratification tool(s) used to assess risk.
 - Lifestyle factors an individual presents with (e.g. inactivity, smoking etc.).
 - The individual's functional movement.
 - The side-effects of medications taken, and number of medications taken.
 - The type and intensity of exercise planned.
 - The exercise environment and factors that affect the environment, e.g. space and temperature.
 - The exercise equipment used.
 - The delivery and instructions skills of the instructor.
 - The ability of the instructor to adapt, modify, and tailor specific exercises.
- A full risk assessment should be undertaken to assess the level of risk.

	• It is the role and responsibility of the exercise instructor to make informed decisions prior to making recommendations for participation in exercise. • Informed decisions need to be based on all the information gathered from participants and this needs to be considered specifically in relation to the type of exercise session and activities being recommended. See 'Appendix 2: Guidance for participant inclusion criteria and scope of practice'.
4.3 Outline the importance of having the correct insurance cover for working with adults with long-term health conditions	• How to check insurance arrangements are sufficient. • Correct insurance requirements for working with adults with long-term health conditions. • Examples of when insurance cover may be insufficient, e.g. depending on qualifications held. • How to inform insurance companies of the specialist qualification and to include cover of this population under existing policy.
4.4 Outline the importance of holding a current and valid first aid and CPR certificate when working with adults with long-term health conditions	• First aid and CPR certification to be completed prior to working with clients with long-term health conditions. • Know how to respond in the event of accidents and/or emergencies. • Recognise signs and symptoms of adverse events. • Provide first aid: ○ Prevent deterioration. ○ Preserve life. ○ Promote recovery.
4.5 Explain the importance of regular continuous professional development (CPD) relevant to working with adults with long-term health conditions	• Maintaining professional competence and membership of organisations. • Keeping up to date with guidance and recommendations. • Best practice and working within scope of practice. • CPD opportunities to support work with this population, including specialist organisations and training providers. • Types of CPD – peer mentoring, clinical service supervision, online. • CPD related to communication and assertiveness, behaviour change, motivational interviewing, health policy, health conditions, palliative care. • Strategies for emotional resilience. • Importance of reflective practice.

4.6 Explain the role of exercise referral schemes in supporting adults with long-term health conditions	• History and overview of schemes including reference to any evidence (where available) to support effectiveness, e.g. evaluation reports available for National Exercise Referral Scheme in Wales. • Where and who schemes are typically delivered by (variable and depends on home country and location). • Primarily to support individuals with low to moderate risk health conditions to participate in structured and supported exercise. • Delivered for specified duration (variable and scheme dependent). • The processes and protocols where delegated responsibility is integrated in the delivery of services for people at high(er) risk of developing or living with LTCs. • With consideration to scheme set-up considerations: ○ Adhere to medico-legal requirements (as per National Quality Assurance Framework – NQAF, 2001). ○ Provide specified inclusion and exclusion criteria. ○ Specific consultation and assessment protocols for assessing clients. ○ Schemes are monitored and evaluated. ○ Schemes are funded, so often reduced costs for established timeframe. ○ Qualified instructors. ○ Other personnel – managers, coordinators, administration staff. ○ Clear roles and responsibilities. ○ Key national and local care pathways supporting physical activity for individuals with specified health conditions (see credible information sources for various sources for specific conditions). • Brief overview and awareness raising of related health policy that impacts scheme development: ○ Relevant UK wide, national, and local physical activity health inequalities. ○ Health and social care policy for the promotion of physical activity, the prevention of inactivity and sedentary behaviour. ○ Policy relating to the management of long-term conditions by use of physical activity. ○ Local health strategies and needs assessments that influence the development of local services, including physical activity and sedentary behaviour.
---	---

	○ The wider determinants of health, their impact on health inequalities and on an individual's health, and the use of health profiles to understand local demographics. ○ The importance and value of monitoring and evaluating service outcomes, including knowledge of key evaluation documents, e.g. Public Health England (PHE) Standard Evaluation Framework and the Sport England Monitoring and Evaluation Framework. ○ The various data methodologies and research areas, e.g. qualitative, quantitative, customer feedback, process evaluation, impact and cost data and the instructor's role in the data collection process. ○ The importance of engaging and maintaining strong relationships with key stakeholders including commissioners, senior managers, community providers, service users, and their carers and how this informs service development of a local physical activity programme. ○ Key national and local care pathways that will link to the provision of local physical activity services including: – Non-clinical, e.g. integrated lifestyle services, social prescribing, Making Every Contact Count (MECC). – Clinical, e.g. diabetes prevention, NHS health checks, IAPT and weight management and rehabilitation services including cardiac, cancer, pulmonary, MSK, neurological, falls.
4.7 Explain the stages of the referral process	● Self-referral pathways: ○ Ensuring the needs of target population are taken into account, including safety considerations. ○ Relevant tools that determine the level of risk of an adverse 'event' during exercise, e.g. PARQ+, PreCise tool, ACSM health screening or other relevant condition-specific tools. ○ The Health Commitment Statement (an agreement <u>not</u> a risk stratification tool). ● Referral from health services: ○ Physiotherapists. ○ GPs. ○ Practice nurses. ○ Dietitians. ● Process outline: ○ Individual visits health care professional who clinically assesses and transfers information.

	○ Information is received and checked and individual signposted forwards (or backwards if incomplete records). ○ Initial assessment of individual. ○ The individual is accepted if within scope and meets the inclusion criteria. ● How to set up an agreed self-referral and referral protocol (including inclusion and exclusion criteria) with local professionals and allied services.
4.8 Explain the difference between a referral, self-referral, a recommendation and signposting and role of the instructor in relation to these	● Self-referral – a client gaining access to services by themselves and not involving a clinical referral. ● Referral – the transfer of care for a patient from one clinician or clinic to another by request. ● Social prescribing – sometimes referred to as community referral, is a means of enabling GPs, nurses, and other primary care professionals to refer people to a range of local, non-clinical services. ● Signposting – a recommendation to another service (not a referral). ● Informed consent does not need to be gained to signpost as no personal information is shared. ● Informed consent is required for referrals where information is shared. ● How to use health knowledge to communicate in a credible manner with health, social care and third sector professionals at different levels working within the health and social care system.
4.9 Explain health, safety and welfare requirements relevant to own working role, including safeguarding	● This content is covered in pre-requisite qualifications and must be explained and applied to new working role and context. ● The importance of health and safety: ○ Everyone has responsibility. ○ Duty of care. ○ Negligence and commission. ○ Safety and wellbeing. ○ Professionalism. ● Relevant requirements and legislation, relating to role: ○ Safeguarding. ○ Risk assessment. ○ Managing emergencies and emergency action plans. ○ Reporting procedures – confidentiality, data protection. ○ First aid regulations and procedures.

	○ Reporting of injuries, diseases, and dangerous occurrences regulations (RIDDOR). ● Key principles for working with vulnerable adults: ○ What is meant by safeguarding? ○ The different types of abuse. ○ Possible signs of abuse. ○ Responsibilities and limitations. ○ Procedures to follow to protect oneself from accusations of abuse. ● Organisational procedures and policies: ○ Disclosure and barring (DBS). ○ Safeguarding adults and adults at risk. ○ Responsible person for managing safeguarding issues. ○ Reporting procedures for safeguarding. ● The statutory agencies responsible for safeguarding: ○ When it may be necessary to contact statutory agencies. ○ How to maintain confidentiality of information relating to possible abuse.
4.10 Identify other working roles designed to support health and physical activity in the community	● Other roles may include: ○ Sport development officers. ○ Exercise referral teams. ○ Link workers. ○ Health champions. ○ Health coaches. ○ Health trainers. ○ Community activators. ● How to use health knowledge to communicate in a credible manner with health, social care and third sector professionals at different levels working within the health and social care system.

Programming, adapting and delivering exercise for adults with long-term health conditions (J/651/1819)

Unit aim

To provide the knowledge and skills to programme, adapt, and deliver exercise sessions and programmes for adults with long-term health conditions, within scope of practice.

Learners will know how to:

- consult with individuals and conduct appropriate health screening and assessment.
- evaluate the potential risks and benefits of participation in exercise.
- assess suitability of exercise for participants within scope of practice.
- signpost participants to other professionals, where needed.
- design, adapt, and deliver safe and effective exercise sessions to meet the needs of individuals with long-term health conditions:
 - provide appropriate adaptations and tailoring to meet specific needs.
 - monitor exercise safety and intensity.
 - demonstrate effective communication and instructional skills.
 - engage and fully support a range of participants.

Content

1. Understand how to screen and assess adults with long-term health conditions prior to participation in physical activity and exercise

1.1 Explain the importance of appropriate consultation, pre-activity screening and assessment when working with adults with long-term health conditions

- Screening methods and risk stratification models:
 - ACSM model – algorithm and current PAR-Q+ with follow-on questions and e-PARmed-X (as required).
 - Irwin and Morgan Model.
 - Verbal screening.
 - Other screening tools for specific conditions (where available and as appropriate).
- With consideration to:
 - Advantages and disadvantages of different methods/models.
 - Evidence-base for use (validity and reliability of tools).

	• Use for: ○ Appropriate advice and guidance – deferral, signposting, and referral. ○ Working within scope of practice and role boundaries, competence, confidence, and qualifications. ○ Identifying suitability of sessions and identifying individuals who need specialist support (physiotherapy). ○ Determining session content. ○ Supporting individuals with lifestyle behaviour change.
1.2 Describe appropriate consultation and communication skills to gather information from participants	• How to conduct a consultation – environment, timing, structure, etc. • The importance of rapport and relationship established between instructor and participant(s) during the consultation on engagement and adherence. • The importance of a person-centred empathetic approach that takes account of the wider determinants of health and the impact on an individual's ability to change their behaviour. • The use of motivational interviewing techniques – open questions, active listening, reflective statements, summaries. • Methods of gathering information: ○ Written questionnaires (PAR-Q+, e-PARmed-X and other questionnaires). ○ Informed consent. ○ Referral and transfer information (where relevant). ○ Interviews and oral questions. ○ Health assessments – blood pressure, heart rate, body mass index (BMI), waist circumference (as appropriate): – When to use and when to exclude, e.g. too invasive. ○ Medical information and medications – transferred by GP or other healthcare professional (HCP). ○ Observation and use of questions throughout session. ○ Benefit of functional assessments, such as three-minute walk, sit to stand, timed up and go, single-leg balance, functional reach, internal and external rotation, seated hamstring stretch. <u>NB: learners are not expected to conduct these assessments, just to have awareness of their value and uses. This information may be available from health care professionals and would be part of more specialist qualifications (e.g. postural stability instructor).</u> ○ Verbal screening at start of every session: – Health and wellbeing checks.

	– Changes to health status or any medications. ○ Other questionnaires that can be useful to identify needs: ▪ EQ-5D – to assess overall wellbeing (see appendix for sources). ▪ Specific to conditions, e.g. Oswestry, PHQ-9. • Appropriateness of methods with consideration to client's needs: ○ Deaf or partial hearing. ○ Blind or partial sighted. ○ Physical disability. ○ Speakers of other languages. ○ Neurodiversity (ADHD, dyspraxia, etc.). • Uses of information: ○ To assist risk stratification. ○ To identify reasons for referral/signposting. ○ To inform planning, delivery, and instruction skills.
1.3 Summarise the information that should be obtained when consulting with and pre-screening adults with long-term health conditions prior to physical activity and exercise	• Information: ○ Personal information – age, gender, etc. ○ Medical and surgical history and medications. ○ Physical activity history and preferences, current fitness. ○ Lifestyle behaviours. ○ Motivation and barriers. ○ Readiness to make changes and goals. ○ Health and physical assessments (where appropriate). ○ Medications and awareness of their side effects, including effects that may impact exercise response, e.g. blood pressure, balance, energy levels, etc.
1.4 Explain when to signpost or refer participants to other healthcare professionals prior to participation in exercise sessions	• When written consent should be sought from healthcare providers: ○ Positive response to PAR-Q+ and follow-on questions, including completion of e-PARmed-X. ○ To gather more information about participants, such as outcomes of any functional or other assessments completed by a health care professional.

	• Risk stratification beyond scope of practice and qualifications: ○ The level of risk and needs identified exceed professional competence/qualifications and type of session being offered. • Issues outside of scope such as: ○ Medical ○ Nutritional ○ Psychological ○ Risk stratification ○ Contraindications. • The importance of participant consent before exchanging information with other professionals. • Other professionals to include: ○ Scheme manager/coordinator ○ GP ○ Counsellor ○ Dietician ○ Smoking cessation ○ Other instructors.
1.5 Outline potential goals of adults with long-term health conditions and the importance of regular participation in physical activity and exercise to support these goals	• Specificity and diversity of goals: ○ Medical management and physiological. ○ General health and fitness and lifestyle. ○ Psychological and social. ○ Psychological. ○ Functional ability. • The importance of person-centred goal setting. • Process, behavioural and outcome goals. • How goals can support motivation. • The effect of goals on session and programme planning and delivery. • Reviewing and checking goals and monitoring progress. • Awareness and overview of when to involve others in goal setting: ○ To maintain scope of practice and role boundaries. ○ Issues outside of scope: – Medical – Nutritional

	– Psychological – Risk stratification – Contraindications. ○ Other professionals to include: – GP – Health trainers – Counsellor – Dietician – Smoking cessation – Other instructors. • The importance of regular participation in physical activity and exercise to support achievement of goals: ○ Physical activity – CMO guidance: – Activities of daily living and messages: ▪ sit down less and move more often. ▪ Some is good, more is better. ▪ Break up sedentary times. ○ Structured exercise: – Frequency, intensity, time, type, volume, and progressions (FITT-VP). – Evidence-based recommendations. – Linked home-based exercise programmes.
1.6 Describe how to record and store information	• Procedures for collecting, storing, and sharing personal and sensitive data that includes medical information. • Records to be maintained with consideration to local GP and NHS or other referral agency information governance requirements, including information imparted by clinicians and/or the participants. • With consideration to: ○ Principles of confidentiality and data protection. ○ General Data Protection Regulations (GDPR). ○ Data Protection Act (2018). ○ The use and value of appropriate electronic data management systems (including local customer relationship management – CRM system, where available):

	– To keep service user records up to date and use this to manage behaviour change and tailor follow up motivational support according to individual needs. – To aid data collection, evaluation, and support to individual participants.
--	---

2. Understand how to plan and adapt exercise for adults with long-term health conditions

2.1 Describe planning considerations for delivering exercise sessions for adults with long-term health conditions	• Consideration to, and/or use of local clinical governance guidelines when delivering physical activity for people at risk of, or living with, an LTC. • Participants: ○ Screening and informed consent. ○ Appropriate clothing and footwear. ○ Hydration. ○ Functional limitations and impairments. ○ Chronic health conditions. ○ Medications and effects. • Environment: ○ Type of environment, e.g. studio, gym, home-based, etc. ○ Temperature, space, floor surface, room layout. • Equipment: ○ Suitability of different equipment for: – Specific populations. – Specific types of session. ○ Maintenance of equipment. ○ Lifting and handling. ○ Types of equipment: – Small portable equipment – resistance bands, chairs, hand weights/dumbbells, ankle weights, steps. – Fixed equipment – gym-based (CV machines, RT machines).
2.2 Describe how to assess and manage the risks in the exercise environment	• Screening and risk stratification of participants: ○ Exercise recommendations and modifications. ○ Referral or deferral as needed. • Effects of chronic health conditions – see Unit 1. • Effects of specific medication(s) – see Unit 1.

	• Risk assessment of environment and equipment: ○ Layout of environment. ○ Equipment used and layout. ○ Including environments not designed for exercise, e.g. individual's home. • Checks needed: ○ Prior to physical activity. ○ During physical activity – dynamic risk assessment, observation. ○ Post physical activity.
2.3 Describe how the effects of long-term health conditions on the body systems may have implications for planning and delivering exercise	• All identified effects and risks – see Unit 1. • Motor skills – balance, coordination, reaction time to instruction and cues. • Pelvic floor – inclusion of specific exercises to target this area. • Transitions: ○ Between movements. ○ From standing to floor and floor to standing and individual's ability and confidence to transition. ○ Planning and allowing time for transitions. ○ Clearer instructions to manage transitions. ○ Use of teaching position to manage transitions. • Slower movement speed (including music speed, where appropriate). • Intensity variables – repetitions, resistance, range of motion, rate/speed, sets. • Modifications for functional impairments: ○ Joint and mobility. ○ Cognitive. ○ Cardiovascular. ○ Sensory – vision and hearing.
2.4 Explain how to apply knowledge of physical activity, ageing and disease processes to design and deliver a safe and effective physical activity programme	With consideration to changes to the body systems relevant to specific health conditions (progressive and over time) include: • Skeletal, i.e. bone density, joint mobility, range of motion (ROM), alignment and posture. • Neuromuscular, i.e. posture, pelvic floor, power, strength, endurance, sarcopenia, heat production, immune function, fine motor control, coordination, balance and falls risk, reaction time (including ability to respond to instructions), movement speed, sight and hearing, short-term memory

	• Cardiovascular, i.e. reduced anaerobic threshold, reduced exercise tolerance, and ability to sustain activity • Respiratory, i.e. breathlessness • Cognitive, i.e. short-term memory • Psychosocial, i.e. low mood, low energy, loss of interest in life
2.5 Describe a safe and effective session structure and appropriate activities for specific long-term health conditions	Session structure considerations, such as: • Warm-up. • Cardiovascular • Muscular • Cooldown and stretch Activities/exercises: • Evidence-based guidelines appropriate to condition. • All components of fitness: ○ Cardiovascular fitness ○ Muscular fitness – functional, for bone density, posture, and pelvic floor ○ Flexibility ○ Motor skills, e.g. balance, coordination, reaction time, speed, etc ○ Relaxation • Activities of daily living (ADL): ○ Move more often and sit down less Types of session • 1:1 – Sessions and programmes • Group-settings – sessions and programmes
2.6 Describe the structure, intensity and exercise selection for a safe and effective warm-up	• For clients with specific long-term health conditions and according to their needs. • Warm-up – longer duration, more progressive and gradual increase of intensity, more targeted and isolated joint mobility for specific joints (where appropriate), lower intensity and impact, transitions, and speed manageable. • Content appropriate to exercise genre.

2.7 Describe the structure, intensity and exercise selection for safe and effective cardiovascular training	• For clients with specific long-term health conditions and according to their needs. • Cardiovascular – lower impact, less intense, progressive, and gradual build-up and cooldown, interval approaches, during peak slower pace and less complex and less arm movements (if included) and less complex transitions. • Content appropriate to exercise genre.
2.8 Describe the structure, intensity and exercise selection for safe and effective muscular fitness training	• For clients with specific long-term health conditions and according to their needs. • Muscular – whole body approach, functional focus and related to daily activities, fracture site focus (osteoporosis), full range of motion and of sufficient intensity to be effective, avoidance of isometric exercises. • Content appropriate to exercise genre.
2.9 Describe the structure, intensity and exercise selection for a safe and effective cooldown	• For clients with specific long-term health conditions and according to their needs. • Cooldown and stretch – stretch positions appropriate, support and balance available (walls, chairs), use of aids to assist range of motion or comfort (pillows, cushions). • Content appropriate to exercise genre.
2.10 Explain the benefits and limitations of different methods of monitoring exercise	• The role of evidence-based technologies that support the uptake and maintenance of physical activity. • Relevant methods of monitoring an individual or group of participants both before, during and after exercise: ○ Rate of perceived exertion – RPE. ○ Talk test. ○ Breathlessness scales. ○ Heart rates / heart rate monitoring. ○ Observation. ○ Questioning. ○ Blood sugar level checks for individuals with diabetes.

2.11 Identify a range of modifications, adaptations and progressions that may be applied to ensure the exercise session and programme is accessible and inclusive to adults with long-term health conditions	• Modification, adaptation, and progression of: ○ Frequency, intensity, time, and type. ○ Intensity variables (to affect modification) and prioritisation of them for increasing/decreasing intensity and/or functional movement: – Range of motion. – Repetitions. – Resistance. – Rate or speed of movement. – Impact. – Exercise positions and start positions. ○ Weight-bearing. ○ Balance and level of support. ○ Music speed – appropriateness of speed and when not appropriate. • Adaptations according to specific condition and level of decline (as appropriate): ○ Joint impairments – shoulder, hip, spine, finger, toes, knees, wrist: – Pain, reduced range of motion (ROM), stiffness, lack of strength, alignment, and posture. ○ Cardiorespiratory – reduced tolerance and capacity, lower muscle mass, breathlessness. ○ Sensory impairment – visual, vestibular, hearing, proprioception. ○ Cognitive/mental – mild dementia, low mood, mild depression/anxiety, decline in cognitive ability, short-term memory, decline in ability to follow instructions safely, decline in ability to sequence tasks.
2.12 Analyse the suitability of a range of alternative types of activity to meet the diverse needs and requirements of adults with long-term health conditions	• Types of activity and who they may be suitable or unsuitable for: ○ Walking programmes (including the risk assessment of some outdoor environments, e.g. uneven walking surface may increase risk of falls). ○ Chair-based exercise. ○ Water-based exercise. ○ Tai Chi. ○ Yoga. ○ Pilates.

	○ Specialist sessions: – Exercise referral. – Cardiac rehabilitation sessions. – Pulmonary rehabilitation. – Strength and balance (postural stability instructor). • When to recommend these alternatives: ○ To maintain scope of practice and competence. ○ Safety and effectiveness for individuals. • When to regress physical activities. • Components of fitness trained or not trained by specific exercise modalities and how to advise participants.
--	---

3. Understand how to deliver and review exercise for adults with long-term health conditions

3.1 Describe how to utilise and adapt communication and instructional skills to monitor and improve performance	• Adaptation according to individual needs and health condition. • The importance of correct technique during physical activity and exercise. • Instructor skills: ○ Accurate own demonstration, including movement speed and posture. ○ Clarity of instruction and cues to support transitions between exercises: – Visual cues. – Verbal cues and use of body language. – Use of voice – volume, intonation, projection, enunciation. – Step-by-step instructions. ○ Timing: – Planning time for transitions. – Time to get into and out of position. – Set-up of start positions. – Time to set up equipment, e.g. use of resistance bands. ○ Observation and use of effective teaching position. ○ Presentation – including clothing, footwear. • The importance of rapport and relationship established between instructor and participants on engagement and adherence.
--	---

3.2 Explain the importance of correct instructions and demonstration of safe and effective exercise technique	• To ensure safe and effective participation.
3.3 Describe how to tailor delivery method to meet the needs of a range of participants	• Adaptation of verbal and instruction methods. • With consideration to learning styles and specific needs, such as: ○ Deaf or partial hearing. ○ Blind or partial sighted. ○ Physical disability. ○ Speakers of other languages. ○ Neurodiversity (ADHD, dyspraxia, etc.).
3.4 Describe effective motivational strategies and techniques to support participants to adhere to physical activity and support lifestyle behaviour(s) change	• Consideration to behaviour change models: ○ Transtheoretical model (stages and processes). ○ Self-efficacy. ○ COM-B (capability, opportunity, motivation, behaviour). ○ Self-determination theory. ○ Health belief model. ○ Biopsychosocial model. • Effective motivational strategies and techniques that can be used to support an individual: ○ Techniques: – Motivational interviewing (OARs and change/sustain talk). – Solution focused. – Cognitive behavioural therapy (CBT). ○ Strategies: – Awareness of self-talk. – Awareness of antecedents and behaviour triggers. – Goal setting and use of rewards. – Social support systems. – Education and learning. – Role models. – Peer support.

3.5 Outline how to apply relevant behaviour change theory and techniques to design a programme to meet the needs of individuals	• See 3.4 • At different stages of intervention: – consultation. – Delivery. – Review points.
3.6 Explain the importance of regular reviews of the participant's progress	• To confirm continued suitability of sessions and programme – safety and effectiveness. • Awareness of need for progression and regression. • To inform health professionals/commissioners. • To evidence statistics and outcomes to gain access to bursaries/grants.
3.7 Identify opportunities to collect feedback from participants	• Before, during, and after sessions. • Information needed – extent to which session met needs, individuals found it difficult or easy, etc. • Methods – verbal, written, reassessment.
3.8 Explain how to use the information gathered from participant feedback to promote motivation, adherence and outcome success	• Review and adapt exercise programmes and behavioural strategies based on individual's biopsychosocial needs and with consideration to: ○ Safety. ○ Motivation levels. ○ Medical information. ○ Personal motivations and preferences. • For example, comparing reports of subjective experiences with outcomes of any functional assessment outcomes/reviews to show progress and developments.
3.9 Explain how to reflect on own practice to inform future sessions	• Kolb Model. • Importance of reflective practice. • Reflection on action and in action. • Use of reflective practice to identify CPD needs.

4. Be able to collect and use information to plan and adapt exercise for adults with long-term health conditions

4.1 Use appropriate methods to collect and record information from participants

- Use of:
 - Clinical information shared by a health care referral:
 - How to interpret this.
 - When and how to refer to referring practitioners.
 - Written (PAR-Q+/e-PARmed-X) and verbal screening.
 - Relevant assessment(s), and risk stratification:
 - Signposting, referral, or deferral – as appropriate.
 - Adaptation of session or content.
 - Progression and regression of session content.
- Different ways to collect information and consult with individuals in different ways:
 - How to develop rapport via face to face, telephone, remote, online approaches, etc.
 - Use a person-centred approach.
 - Integrate the use of behaviour change models and motivational techniques.
 - The pros and cons of different methodologies, the evidence for these and impact on practice in selection and use and understand how to use within a person-centred behaviour change approach.
- Readiness of individual to change lifestyle behaviours (motivation, confidence, stage of behaviour change) all of which will dictate the amount of support needed and prioritisation in session.
- When assessments are used:
 - Educate client on purpose and value.
 - Select assessments appropriate to the individuals.
 - Advise individuals of correct procedures, protocols, and risks prior to commencing any physical assessment(s).
 - Gain informed consent prior to completion of any assessment.
 - Supervise assessments in a safe and effective manner.
 - Carry out assessments at an appropriate point in a behaviour change intervention, in an empathetic and non-judgemental style.

	○ Ensure safety of client and professionalism is maintained at all times: – Correct procedures, protocols. – The presence of a chaperone, if necessary. – Informed consent. – Duty of care. – Cultural sensitivities.
4.2 Analyse and use information gathered to design a safe and effective exercise session for adults with specific long-term health conditions	● Interpret and use the results of information gathered to: ○ Establish a base line from which to monitor and review progress and outcomes and adapt programmes at regular intervals as appropriate. ○ Make appropriate lifestyle recommendations (within scope of practice). ○ Signpost individuals with needs that exceed scope. ○ Design exercise programmes based on individual's biopsychosocial needs and with consideration to: – Clinical information and needs. – Safety. – Motivation levels. – Medical information. – Personal motivations and preferences. ○ Safe and effective session structure: – Warm-up. – Cardiovascular exercises. – Muscular fitness exercises. – Cooldown and flexibility exercises – relaxation. – With consideration to: ▪ Motor skills, including balance, coordination. ▪ Functional considerations – ability and confidence to transition. ▪ Recommendations for other activities – activities of daily living, home-based exercise, other exercise sessions appropriate to needs. ● Application of progressive principles and variables (FITT-VP).

4.3 Apply relevant behaviour change theory and techniques to design a programme to meet the personal needs of individuals	• See 3.4 and 3.5 • Application of principles from relevant behaviour change models: ○ Transtheoretical model (stages and processes) ○ Self-efficacy ○ COM-B (capability, opportunity, motivation, behaviour) ○ Self-determination theory ○ Health belief model ○ Biopsychosocial model
4.4 Plan a safe and effective warm-up component	• Intensity, duration, and selection of exercises to meet the needs of participants. • Content appropriate to exercise genre.
4.5 Plan a safe and effective main component	• Intensity, duration, and selection of exercises to meet the needs of participants. • Content appropriate to exercise genre.
4.6 Plan a safe and effective cooldown and stretch component	• Intensity, duration, and selection of exercises to meet the needs of participants. • Content appropriate to exercise genre.
4.7 Assess and manage risks in the exercise environment	• Individual risks – conditions, side effects of medications, reliever medications available (respiratory conditions), carbohydrate snacks available (diabetes), etc. • Environment risks – temperature, etc. • Equipment risks – safe lifting and moving. • Management of risks (eliminate, reduce, isolate, control) and contingency plans.
4.8 Provide a rationale for the session structure and activities used in the session/programme	• Purpose of session. • Reasons for activities selected according to participant(s) needs. • Reasons for the exclusion of any components, related to genre delivered and advice that would be provided to participants regarding training other components.
4.9 Plan a range of adaptations, modifications and progressions for the exercise programme specific to participants' needs	• Consider that health conditions and individual health status can change over-time. • Application of FITT-VP principles: ○ Frequency ○ Intensity: – Repetitions

	– Rate/speed – Resistance – Rest – Range of movement. ○ Time: – Whole session – Components. ○ Type: – Modify or change exercise modality or exercises – Impact and weightbearing changes. ○ Volume – dose-response ○ Progression – according to individual needs. • Different training approaches for component of fitness and exercise discipline (appropriate to type of session and genre planned): ○ Cardiovascular: continuous, intervals, etc. ○ Resistance: circuit, single sets, multiple sets, etc. ○ Flexibility: assisted, dynamic, static (modified and adapted). • Advice to provide to individuals for sessions where certain components of fitness are not trained.
4.10 Record the session and programme in an appropriate format	• Records to be maintained with consideration to local and NHS or other referral agency information governance requirements and information imparted by clinicians and/or the participants. • For assessment, use records provided in the Learner Assessment Record (LAR). • The importance of accurate records when working with individuals with long-term health conditions. • Use of appropriate screening records, risk assessment. • Use of appropriate planning records – lesson plan and progressive plan. • Records to be maintained for: ○ Monitoring and reporting purposes (as appropriate). ○ Protection in the event of litigation. • Records to align with GDPR and data protection guidelines: ○ Essential information. ○ Confidentiality. ○ Clear and structured.

	○ Appropriate format. ○ Appropriate storage and transfer.
--	--

5. Be able to deliver and review exercise sessions and programmes for adults with long-term health conditions

5.1 Pre-screen participants prior to participation to check readiness to participate	• See 3.1 and 3.3
5.2 Assess, monitor and manage risk to participants throughout the programme	• See 3.2
5.3 Deliver a safe and effective warm-up component	• Use of appropriate instructional skills. • Intensity, duration, and selection of exercises to meet the needs of participants.
5.4 Deliver a safe and effective main component	• Use of appropriate instructional skills. • Intensity, duration, and selection of exercises to meet the needs of participants.
5.5 Deliver a safe and effective cooldown and stretch component	• Use of appropriate instructional skills. • Intensity, duration, and selection of exercises to meet the needs of participants.
5.6 Use appropriate methods of monitoring exercise safety and intensity	• Methods of monitoring exercise safety and intensity appropriate to health conditions and effects of medication on the exercise response: ○ Observation ○ Questions and answer ○ Heart rate ○ Rating of perceived exertion (RPE) ○ Talk test
5.7 Observe and correct participants' exercise technique to ensure safe and effective alignment and use of equipment where appropriate	• Change of teaching position. • Use of eye contact. • Use of specific teaching points.

5.8 Use effective communication and instructional skills and a person-centred and empathic approach to deliver the session	• Adopt a person-centred empathetic approach that takes account of the wider determinants of health and the impact on an individual's ability to change their behaviour. • Application of all the following with consideration to the individual's physical, psychological, and behavioural needs: ○ Visual and verbal communication skills ○ Teaching points ○ Demonstrations ○ Explanations ○ Teaching position ○ Alternatives ○ Observation and correction ○ Motivational strategies. • Adaptation of different methods of communication for specific conditions/clients (as required).
5.9 Provide client specific instructing points, feedback and reinforcement in a friendly, professional manner	• Based on participant's performance and specific needs.
5.10 Adapt and tailor delivery method, session structure and activities to meet the needs of all participants	• Regressions of exercises, as appropriate. • Progressions of exercises, as appropriate. • Change of teaching position to improve observation. • Modification of instruction skill.
5.11 Provide adaptations, alternatives and progressions to meet individual needs (as required)	• As required to support participants.
5.12 Use effective motivational strategies and techniques to support participants to adhere to physical activity and support lifestyle behaviour(s) change	• Verbal and visual • General and specific to individuals • Praise • Encouragement

5.13 Evaluate the effectiveness of the session and programme to ensure it is engaging, varied and progressive to participants' needs/goals	• Regular review dates. • Use of participant feedback and self-reflection. • Things that went well – session content and delivery. • Things to change or improve – session content and delivery.
5.14 Use information gathered from participant feedback to promote motivation, adherence and outcome success	• Measures of success may include: – long-term adherence. – changes to activity behaviour – subjective measures, such as improved mood and energy levels
5.15 Use information gathered from participant feedback and self-reflection to inform own continuing professional development	• How to improve own practice. • The value of reflective practice. • Relevant continuing professional development (CPD).

Appendix 1: Information sources

Please note while the information sources listed are available at the point of development/publication, access to specific website pages will change over time, as will the currency of information.

Information sources and organisations

- Age UK: <https://www.ageuk.org.uk/>.
- Age Concern: <https://regenerage.org.uk/>
- American College of Sports Medicine (ACSM): <https://www.acsm.org/>.
- American Diabetes Society (ADA): <https://www.diabetes.org.uk/>.
- Association for Nutrition (AfN): <https://www.associationfornutrition.org/>.
- Asthma UK: <https://www.asthmaandlung.org.uk/>.
- Arthritis action: <https://www.arthritisaction.org.uk>.
- Blood Pressure UK: <https://www.bloodpressureuk.org/>.
- British Association of Cardiovascular Prevention and Rehabilitation (BACPR): <https://www.bacpr.org/>.
- British Journal of Sports Medicine: <https://bjsm.bmj.com/>.
- British Medical Journal (BMJ): <https://www.bmj.com/>.
- British Heart Foundation (BHF): <https://www.bhf.org.uk/>.
- British Geriatric Society: www.bgs.org.uk/.
- Blood Pressure Association: www.bpassoc.org.uk/.
- British Association of Sports and Exercise Science (BASES): <https://www.bases.org.uk/>.
- British Lung Society: <https://www.lunguk.org/>.
- British National Formulary (BNF): <https://about.medicinescomplete.com/#/>.
- British Nutrition Foundation: <https://www.nutrition.org.uk/>.
- Centre for behaviour change: <https://www.ucl.ac.uk/behaviour-change/>.
- Centre for ageing better: <https://ageing-better.org.uk/ageing-population>.
- Chartered Institute for the Management of Sport and Physical Activity (CIMSPA): <https://www.cimspa.co.uk/>.
- Diabetes Education and Self-Management for Ongoing and Newly Diagnosed (Desmond) <https://www.desmond.nhs.uk/>.
- Diabetes UK: <https://www.diabetes.org.uk/>.
- Dose Adjustment for Normal Eating (Dafne): <https://dafne.nhs.uk/>.
- e-PARmed-X + screening tool: <https://eparmedx.com/>.
- EQ-5D assessments: <https://euroqol.org/eq-5d-instruments/>.

- Health and Safety Executive (HSE): <https://www.hse.gov.uk/>.
- Journal of Public Health: <https://academic.oup.com/jpubhealth>.
- Later Life Training: laterlifetraining.co.uk/
- Map of medicine: <https://mapofmedicine.com/>.
- Mental Health Foundation: <https://www.mentalhealth.org.uk/>.
- MIMS: <https://www.mims.co.uk/drugs>.
- MIND: <https://www.mind.org.uk/>.
- Motivational Interviewing:
<https://motivationalinterviewing.org/understanding-motivational-interviewing>.
- MS Society: <https://www.mssociety.org.uk/about-ms/what-is-ms>.
- National Library of Medicine: <https://pubmed.ncbi.nlm.nih.gov/>.
- National Rheumatoid Arthritis Society: <https://nras.org.uk/>.
- National Institute of Mental Health (NIMH): www.nimh.nih.gov.
- National Institute of Health and Care Excellence (NICE): <https://www.nice.org.uk/>.
- NHS Choices: <https://www.nhs.uk/>.
- National Institute of Health and Care Excellence (NICE): www.nice.org.uk/.
- National Library of Medicine: pubmed.ncbi.nlm.nih.gov/.
- PAR-Q+: <https://eparmedx.com/>.
- Parkinson's UK: <https://www.parkinsons.org.uk/information-and-support/what-parkinsons>.
- Patient UK: <https://patient.info/>.
- Psychology of Sport and Exercise:
<https://www.sciencedirect.com/journal/psychology-of-sport-and-exercise>.
- Research Quarterly for Exercise and Sport:
<https://www.tandfonline.com/toc/urqe20/current>.
- Royal Osteoporosis Society: <https://theros.org.uk/>.
- Sport England: <https://www.sportengland.org/>.
- Sport England Market Segmentation: <https://segments.sportengland.org/pdf/quickGuide-clubs.pdf>.
- Scottish Intercollegiate Guidelines Network (SIGN): <https://www.sign.ac.uk/>.
- The King's Fund. <https://www.kingsfund.org.uk>.
- UK Active: <https://www.ukactive.com/>.
- World Health Organization: <https://www.who.int/>.

Reports and articles

- Age UK (2015), '1m more living with multiple conditions by 2020', UK, Accessed on: 25-05-2026. Available at: www.ageuk.org.uk/latest-press/archive/one-million-more-older-people-will-be-living-with-multiple-long-term-conditions/
- Age UK (2015), 'A practical guide to healthy ageing (70+)', UK, Accessed on: 25-05-2026 Available at: <https://www.england.nhs.uk/publication/practical-guide-to-healthy-ageing/>
- Age UK (2019), 'Later life in the United Kingdom', UK, Accessed on: 25-05-2026 Available at: www.ageuk.org.uk/globalassets/age-uk/documents/reports-and-publications/late_life_uk_factsheet.pdf
- Armstrong, M. et al. (2024), 'Where to Start? Physical Assessment, Readiness, and Exercise Recommendations for people with Type 1 or Type 2 Diabetes', US, Accessed on: 25-05-2026. Available at: <https://pubmed.ncbi.nlm.nih.gov/37193205/>
- Barnett, K., Mercer, S. W., Norbury, M., Watt G., Wyke S., Guthrie, B. (2012), 'Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study.' The Lancet online, UK, Accessed on: 25-05-2026 Available at: [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(12\)60240-2/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(12)60240-2/fulltext)
- British Geriatric Society (2019), 'Strong, straight and steady', UK, Accessed on: 25-05-2026 Available at: www.bgs.org.uk/resources/strong-steady-straight-nos-exercise-and-osteoporosis-consensus-statement
- British Heart Foundation National Centre for Physical Activity (BHFNC) (2010) 'A Toolkit for the Design, Implementation and Evaluation of Exercise Referral Schemes'. UK. Accessed on: 25-05-2026 Available at: ncsem-em.org.uk/wp-content/uploads/2020/10/section-6-exercise-referral-scheme-coordinators.pdf
- Centre for Ageing Better (2019), 'Raising the Bar on Strength and Balance', UK, Accessed on: 25-05-2026 Available at: <https://ageing-better.org.uk/resources/raising-bar-strength-balance>
- Centre for Ageing Better (2021), 'Keep on Moving: Understanding Physical Inactivity Among 50–70-year-olds', UK, Accessed on: 25-05-2026 Available at: <https://ageing-better.org.uk/resources/keep-on-moving-understanding-physical-inactivity-among-50-70-year-olds>
- Centre for Ageing Better (2022), 'The State of Ageing', UK, Accessed on: 25-05-2026 Available at: ageing-better.org.uk/state-of-ageing
- Chief Medical Officer (CMO) (2019), 'UK Chief Medical Officers' Physical Activity Guidelines', UK, Accessed on: 25-05-2026 Available at: <https://www.gov.uk/government/publications/physical-activity-guidelines-adults-and-older-adults>
- Department of Health (2001) Exercise Referral Systems: A National Quality Assurance Framework. London: Department of Health. Accessed on: 25-05-2026 Available at: http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_4009671
- Hall, R., The Guardian (2021) 'A third of middle-aged UK adults have at least two chronic health conditions'. UK, Accessed on: 25-05-2026 Available at:

www.theguardian.com/society/2021/jul/28/a-third-of-middle-aged-uk-adults-have-at-least-two-chronic-health-issues-study

- Irwin and Morgan (2010) 'A Toolkit for the Design, Implementation and Evaluation of Exercise Referral Schemes'. UK. Accessed on: 25-05-2026 Available at: ncsem-em.org.uk/wp-content/uploads/2020/10/section-6-exercise-referral-scheme-coordinators.pdf.
- NICE (2014), 'Behaviour Change. Individual Approaches. Public Health Guidance PH49', UK, Accessed on: 25-05-2026. Available at: <https://www.nice.org.uk/guidance/ph49>
- NICE (2014), 'Physical Activity. Exercise Referral Schemes. Public Health Guidance PH54', UK, Accessed on: 25-05-2026. Available at: <https://www.nice.org.uk/guidance/ph54>
- Richmond Group of Charities (2018), 'Multimorbidity. Understanding the Challenge', UK, Accessed on: 25-05-2026. Available at: https://richmondgroupofcharities.org.uk/sites/default/files/multimorbidity_-_understanding_the_challenge.pdf
- Royal Osteoporosis Society (2018), 'Strong, Straight and Steady'. UK, Accessed on: 25-05-2026 Available at: <https://theros.org.uk/clinical-publications-and-resources/>
- Sport England (2022), 'Easier to be Active', UK, Accessed on: 25-05-2026 Available at: www.sportengland.org/news/new-guidance-help-people-long-term-health-conditions-get-active
- Sport England (2022), 'New guidance to help people with long-term health conditions get active', UK, Accessed on: 25-05-2026 Available at: <https://www.sportengland.org/news/new-guidance-help-people-long-term-health-conditions-get-active>
- UK Active (2021), 'Life in Our Years'. UK, Accessed on: 25-05-2026 Available at: <https://www.ukactive.com/reports/life-in-our-years/>
- Zahalka, S., et al. (2024), 'The Role of Exercise in Diabetes', UK, Accessed on: 25-05-2026. Available at: <https://pubmed.ncbi.nlm.nih.gov/31751111/>

Textbooks

- American College of Sports Medicine (2018) ACSM's Guidelines for Exercise Testing and Prescription. 10th edition. USA: Wolters Kluwer
- American College of Sports Medicine (2022) ACSM's Guidelines for Exercise Testing and Prescription. 11th edition. USA: Wolters Kluwer
- Durstine, L. J., et al. (2009) ACSM's Exercise Management for Persons with Chronic Diseases and Disabilities. USA: Human Kinetics
- Lawrence, D. (2013) The Complete Guide to Exercise Referral. UK: Bloomsbury Publishing
- Scott, A., and Broom, D. (Eds.) (2022) Exercise Management for Referred Medical Conditions. UK: Routledge

Appendix 2: Guidance for participant inclusion criteria and scope of practice

The following screening tools can be used as a **guide** to help identify individuals whose needs are within scope of practice.

Using ACSM algorithm

- Inactive individuals with:
 - No CV, metabolic or renal disease – no medical clearance needed.
 - No signs and symptoms of CV, metabolic or renal disease – no medical clearance needed.
 - Known CV, metabolic or renal disease (asymptomatic) – medical clearance recommended.
 - Signs and symptoms of CV, metabolic or renal disease – medical clearance recommended.
- Active individuals with:
 - No CV, metabolic or renal disease – no medical clearance needed.
 - No signs and symptoms of CV, metabolic or renal disease – no medical clearance needed.
 - Known CV, metabolic or renal disease (asymptomatic) – medical clearance not needed for moderate intensity activity.
 - Signs and symptoms of CV, metabolic or renal disease – seek medical clearance.

NB: Medical clearance being provided does not mean an individual's needs are within your scope of practice unless additional qualifications are held.

Using PAR-Q+ and ePARmed-X

- 'No' response to PAR-Q+ – **no medical clearance needed**.
- 'Yes' response to PAR-Q+ but 'no' response to follow on questions – **no medical clearance needed but participants may require significant modification to some activities (intensity, type, duration)**.
- 'Yes' response to PAR-Q+ and 'yes' response to follow on questions – completion of e-PARmed-X recommended available at www.eparmedx.com and medical clearance may be needed.

Using Irwin and Morgan

- Low to moderate risk stratification

Appendix 3: Conditions within scope of practice for this qualification

These are offered as a guide for inclusion and exclusion criteria. Learners are required to undertake risk stratification to assess the suitability of exercise for a client's needs.

*** NB: If other CVD risk factors exist, or multiple health conditions are present, risk stratification will increase and may exceed scope of practice.**

Condition	Inclusion criteria
Hypertension	Stable and controlled*
Hypercholesterolaemia	Stable and controlled*
Hyperlipidaemia	Stable and controlled*
Hypothyroidism	Stable and controlled*
Hyperthyroidism	Stable and controlled*
Coronary heart disease	Post-phase IV rehabilitation. No new referrals. CVD risk factors in moderate category. No MI.
Peripheral vascular disease	Without other complications
Diabetes type 1 and 2	Stable and controlled with no complications * <i>HCP to advise on modification of insulin prior to exercise (if insulin dependent)</i>
Obesity	Body Mass Index (BMI) 30 - <40 *
Chronic obstructive pulmonary disease (COPD)	Mild/moderate only *
Asthma	Stable and well controlled
Osteoarthritis	Level 1 and 2 only
Rheumatoid arthritis	Early stage and moderate only. Not in flare up.
Osteopenia	Be able to walk independently, no history of hip or vertebrae fractures within the last 12 months

Condition	Inclusion criteria
Osteoporosis	Early stage and moderate. No fracture history. No falls risk
Low back pain	Non-specific only No red flags. Yellow (psychological) flags may require signposting to pain management
Joint replacement	Post-rehabilitation/physiotherapy
Depression	Less severe (mild to moderate only)
Stress	Free from absolute contraindications. Any identified Relative contraindications are stable and controlled.
General anxiety disorder (GAD)	Less severe (mild to moderate)

Please note:

- When multiple health conditions are present, this may increase risk stratification of individuals.
- Individuals with higher or high-risk stratification are outside of scope of practice for this qualification.
- Medical clearance does not mean that all types of exercise are appropriate.
- Instructors should use a range of risk stratification tools to assess level of risk and make informed decisions regarding the specificity of risk, (cardiac incident, fall, fracture, sprain, hypoglycaemia, etc.), severity and likelihood of risk and professional scope of practice.

Exclusions – low back pain (red flags)

- Non-mechanical pain, e.g. nerve root pain
- Thoracic pain
- Fever and unexplained weight loss
- Bladder or bowel dysfunction
- History of carcinoma (cancer)
- Ill-health or presence of other illness
- HIV
- Progressive neurological deficit
- Disturbed gait or saddle anaesthesia (cauda equina)
- Age of onset 55 years

Exclusions – absolute contraindications (exercise referral toolkit 2010)

- BMI <18.5 or >40kg/m²
- Symptomatic severe aortic stenosis
- Acute pulmonary embolus or pulmonary infarction
- Acute myocarditis or pericarditis
- Suspected or known dissecting aneurysm
- Active retinal haemorrhage
- Resting systolic blood pressure ≥ 180mmHg/diastolic blood pressure ≥ 100mmHg
- Uncontrolled/unstable angina
- Acute uncontrolled psychiatric illness
- Unstable or acute heart failure
- New or uncontrolled arrhythmias
- Other rapidly progressing terminal illness
- Significant drop in BP during exercise
- Uncontrolled resting tachycardia ≥ 100 bpm
- Febrile illness
- Experiences pain, dizziness, or excessive breathlessness during exertion
- Unstable/uncontrolled diabetes
- Unstable/uncontrolled cardiac disease
- Severe rheumatoid and osteoarthritis
- Any other unstable, uncontrolled condition
- Any conditions not covered in this qualification

Appendix A: Signs, symptoms, pathophysiology and causes

Hypertension

Signs and symptoms	Pathophysiology	Risks and causes
Asymptomatic; nosebleeds; shortness of breath; general malaise; flushed appearance	Sustained elevated blood pressure >140/90 mmHg.	At risk for heart disease, heart attack, stroke/TIA, heart failure, peripheral arterial disease, aortic aneurysm, kidney disease and vascular dementia. Modifiable risk factors: obesity, sedentary lifestyle, high salt intake, smoking, excess alcohol, excess caffeine, stress, socioeconomic factors e.g. living in a deprived area. Non-modifiable risk factors: age >65 years, family history, genetics, Black African/Black Caribbean ethnicity or descendent of. Known causes: kidney disease, diabetes, sleep apnoea, hormone disorders, lupus, scleroderma.

Hyperlipidaemia

Signs and symptoms	Pathophysiology	Risks and causes
Usually asymptomatic	Elevated cholesterol and triglycerides.	At risk for atherosclerosis, arteriosclerosis, heart attack, stroke and peripheral artery disease. Modifiable risk factors: high saturated fat diet, sedentary lifestyle, obesity. Non-modifiable risk factors: having other conditions e.g. diabetes, menopause, genetics: familial hyperlipidaemia.

Hypercholesterolaemia

Signs and symptoms	Pathophysiology	Risks and causes
Usually asymptomatic	Blood cholesterol levels >5 mmol/L; plaque/atheroma build-up in artery walls.	At risk for atherosclerosis, arteriosclerosis, angina, hypertension and stroke. Modifiable risk factors: saturated fat intake, obesity, smoking, alcohol, sedentary lifestyle. Non-modifiable factors: age, ethnicity, family history, gender, menopause.

Coronary Heart Disease

Signs and symptoms	Pathophysiology	Risks and causes
Chest pain; shortness of breath; pain in neck, shoulders, jaw or arms; feeling faint; nausea	Plaque/atheroma build-up within coronary arteries (atherosclerosis) leads to reducing or blockage of blood supply to heart.	At risk for angina, heart attack and heart failure . Modifiable risk factors: smoking, hypertension, hypercholesterolaemia, hyperlipidaemia, sedentary lifestyle, obesity. Non-modifiable risk factors: family history, diabetes, familial hyperlipidaemia.

Peripheral Vascular Disease

Signs and symptoms	Pathophysiology	Risks and causes
Can be asymptomatic. Dull leg pain or cramping on exertion relieved by rest; numbness or weakness in legs; persistent ulcers on legs and feet; hair loss on legs and feet.	Plaque/atheroma build-up in leg arteries(atherosclerosis).	At risk of progressing to critical limb ischaemia (CLI) and coronary heart disease (CHD). Modifiable risk factors: smoking, hypertension, hypercholesterolaemia. Non-modifiable risk factors: diabetes, age.

Obesity

Signs and symptoms	Pathophysiology	Risks and causes
BMI >30 kg/m ² (or >27.5 kg/m ² in Asian, Chinese, Middle Eastern, Black African or African Caribbean descent); waist-to-height ratio >0.5	Excess body fat.	There is no single conclusive cause of obesity; it is a multifactorial condition influenced by genetic, behavioural, environmental and socioeconomic factors At risk for cardiovascular disease (CVD), respiratory disease, cancers, Type 2 diabetes, joint problems and mental health conditions. Modifiable risk factor: excess calorie intake, poor diet, sedentary lifestyle, socioeconomic influence (fast food, reliance on technology). Non-modifiable risk factors: genetics, medical conditions such as hypothyroidism and Cushing's syndrome

Type 1 Diabetes

Signs and symptoms	Pathophysiology	Risks and causes
Excessive thirst; frequent urination; fatigue; unintentional weight loss; blurred vision; slow wound healing; fruity-smelling breath	Autoimmune disease which attacks the cells in the pancreas, which is responsible for the production of insulin, resulting in insulin deficiency; abnormally high blood glucose levels (hyperglycaemia).	At risk of hypoglycaemia (blood glucose levels below 4mmol/l) and hyperglycaemia (blood glucose levels above 10mmol/l). May progress to peripheral neuropathy, retinopathy, silent angina/MI; diabetic kidney disease, neuropathy cardiovascular diseases (CVD), diabetic ketoacidosis (DKA), other immune diseases; anxiety and depression. Environmental factors: obesity, viruses, vitamin D exposure, diet. Non-modifiable risk factors: viruses, genetics and family history.

Type 2 Diabetes

Signs and symptoms	Pathophysiology	Risks and causes
Can be asymptomatic; excessive thirst; frequent urination; fatigue; unintentional weight loss; blurred vision; slow wound healing	Abnormally high blood glucose levels (hyperglycaemia); insufficient insulin production and reduced insulin sensitivity	At risk of hypoglycaemia (blood glucose levels below 4mmol/l) and hyperglycaemia (blood glucose levels above 10mmol/l). And potential to progress to cardiovascular disease (CVD), dyslipidaemia; diabetic kidney disease, peripheral neuropathy, retinopathy, diabetic foot complications, diabetic ketoacidosis (DKA), anxiety and depression. Modifiable risk factors: obesity, sedentary lifestyle, low-fibre/high-GI diet. Non-modifiable risk factors: ethnicity (people of Asian, African and Afro-Caribbean descent), family history of gestational diabetes, polycystic ovary syndrome (PCOS), certain medications, low birth weight.

Hyperthyroidism

Signs and symptoms	Pathophysiology	Risks and causes
Anxiety; palpitations; twitching or tremor; weight loss; diarrhoea; heat sensitivity; persistent tiredness and weakness; agitation; enlarged thyroid gland (goitre)	Excess thyroid hormone production	Commonly due to Graves' disease or nodules on thyroid gland (non-cancerous). Less commonly due to swelling of the thyroid (thyroiditis), thyroid cancer and pituitary adenoma. Untreated disease may lead to underactive thyroid; osteoporosis, pregnancy complications (pre-eclampsia, miscarriage and premature labour) heart failure, thyroid storm and thyroid eye disease. Modifiable risk factors: smoking, excessive iodine intake. Non-modifiable risk factors: gender (females at greater risk than males), family history, other autoimmune disease.

Hypothyroidism

Signs and symptoms	Pathophysiology	Risks and causes
Fatigue; weight gain; depression; cold sensitivity; dry skin and hair; muscle aches	Insufficient thyroid hormone production.	The causes of an underactive thyroid include Autoimmune disease (Hashimoto's disease); damage to the thyroid due to treatments for an overactive thyroid or thyroid cancer If untreated at risk of heart problems, goitre, pregnancy complications and myxoedema coma (very rare but life threatening) Modifiable risk factors: Certain medications (if the benefits outweigh the risks). Non-modifiable risk factors: family history of Hashimoto's disease.

Asthma

Signs and symptoms	Pathophysiology	Risks and causes
Wheezing; chest tightness; breathlessness; coughing	Chronic lung condition causing temporary narrowing of the airways due to inflammation, mucus secretion and broncho-constriction, causing difficulty breathing. Reduction in FEV1 (volume of air forceable expired in one second) and reduction in peak expiratory flow rate (PEFR) Asthma can be triggered by allergies; exercise; infection.	Modifiable risk factors: obesity, smoking, living or working in urban areas; exposure to environmental allergens e.g. pollution; occupational exposure to certain chemicals, fumes and dust. Non-modifiable risk factors: family history, allergies, low birthweight, childhood exposure to tobacco smoke and air pollution; childhood respiratory infections.

Chronic Obstructive Pulmonary Disease (COPD)

Signs and symptoms	Pathophysiology	Risks and causes
Chest tightness; shortness of breath on mild exertion; chronic or chesty cough; wheezing	COPD includes long-term (chronic) bronchitis and emphysema, characterised by obstruction to airflow that interferes with normal breathing. • Chronic bronchitis – long-term inflammation and narrowing of the airways. • Emphysema – damage and loss of elasticity in the air sacs (alveoli) in the lungs. Airways are narrowed due to: • Lung tissue damage reducing pull on airways • Mucus secretions blocking part of the airway • Airway lining is inflamed and swollen.	Modifiable risk factors: smoking, exposure to environmental smoke/pollution, occupational exposure to certain chemicals, fumes and dust. Non-modifiable risk factors: genetics, age, viral or bacterial infections; history of childhood respiratory disease; a rare genetic condition called alpha-1 antitrypsin deficiency

Lower back pain

Signs and symptoms	Pathophysiology	Risks and causes
Low back pain; radiating pain to buttocks or thighs; muscle spasm	Acute (<3 months) or chronic (>3 months).	May impair work, sleep, leisure, and increase the risk of falls; chronic pain; anxiety and depression. Modifiable risk factors: poor posture; smoking; obesity, sedentary, occupations that include heavy lifting/bending/twisting, stress or depression. Non-modifiable risk factors: age, genetics, accident/injury, congenital or structural differences (spinal variations)

Osteoarthritis

Signs and symptoms	Pathophysiology	Risks and causes
Pain at rest and exercise; stiffness; swelling; muscle weakness	Long-term disorder where damage to a synovial joint causes the body to attempt repair, but these repair processes gradually alter the structure of the joint • Localised cartilage loss • Remodelling of bones and formation of osteophytes • Mild synovitis (inflammation of the synovial membrane)	At risk of progressing to joint deformity, functional limitations (e.g. grip strength, difficulty walking and climbing stairs), chronic pain and mental health impacts (anxiety, depression, self-confidence, self-esteem etc.) Modifiable risk factors: obesity, sedentary lifestyle, occupations that include repetitive squatting/lifting/bending, some sports and exercise stresses, repetitive strain injury (RSI). Non-modifiable risk factors: age, injury, family history, gender (females at greater at risk than males), joint misalignment.

Rheumatoid arthritis

Signs and symptoms	Pathophysiology	Risks and causes
Pain; swelling; stiffness; heat; limited range of movement (ROM) Additional systemic symptoms can include fatigue; high temperature; sweating; loss of appetite; weight loss	Chronic systemic inflammatory disease primarily affecting small synovial joints in hands, feet and wrists. Can develop to cause inflammation in lungs, heart, eyes and blood vessels.	Causes unknown. Risk factor for joint damage, osteoporosis, cardiovascular disease (CVD), anaemia and infection. Modifiable risk factors: smoking. Non-modifiable factors: family history; gender (females at greater at risk than males)

Osteopenia

Signs and symptoms	Pathophysiology	Risks and causes
Usually asymptomatic	Reduced bone mineral density = BMD -1 to -2.5 SD (standard deviation).	Risk of osteoporosis. Modifiable risk factors: sedentary lifestyle, excess alcohol, amenorrhoea (absence of menstrual periods) due to excessive exercise/dieting. Non-modifiable factor: gender (females at greater at risk than males), early menopause (before age 45), early hysterectomy (before age 45)

Osteoporosis

Signs and symptoms	Pathophysiology	Risks and causes
Usually asymptomatic Bone fractures on slight impact; kyphosis; loss of height; stooped posture due to osteoporosis in the spine.	Reduced bone mineral density below -2.5 SD.	At risk of increased fractures and falls. Likelihood of requiring hip replacements Modifiable risk factors: sedentary lifestyle, excess alcohol, smoking, amenorrhoea (absence of menstrual periods) due to excessive exercise/dieting; low BMI <19, steroid use. Non-modifiable risk factors: gender (females at greater at risk than males), family history, eating disorder history, certain conditions e.g. Coeliac disease and Crohn's disease; early menopause (before age 45), early hysterectomy (before age 45)

Joint replacement

Signs and symptoms	Pathophysiology	Risks and causes
Significant pain; inability to perform daily tasks without aid; advanced osteoporosis; significant joint damage	Partial or total joint replacement With post-surgical rehabilitation normal function is expected within 6 months	Risk of dislocation (new joints prone to dislocation post operation due to instability) and future replacement (lifespan of a joint replacement is 10 years). Modifiable risk factors: chronic obesity. Non-modifiable risk factors: osteoarthritis, injury, trauma.

Depression

Signs and symptoms	Pathophysiology	Risks and causes
Persistent low mood; loss of interest in activities; sleep changes; weight changes; appetite changes; fatigue; agitation; poor concentration; indecisiveness; worthlessness; guilt; suicidal thoughts	Reduced positive affect with emotional, physical, behavioural and cognitive symptoms for at least 2 weeks. Defined as: • Less severe depression (subthreshold and mild depression) • More severe depression (moderate and severe depression)	At risk of mental health issues could impair daily functioning and worsen symptoms of co-existing diseases; alcohol and substance misuse; self-harm; self-neglect; suicide Environmental risk factors: unemployment, homelessness, poverty, bereavement, social isolation. Non-modifiable risk factors: family history, gender, menopause, pregnancy, giving birth history of depression or other mental health conditions, substance misuse history; other chronic physical health conditions.

Stress

Signs and symptoms	Pathophysiology	Risks and causes
Headaches; dizziness; muscle tension or pain; elevated heart rate; chest pain; poor concentration; indecisiveness; feeling overwhelmed; constant worrying; forgetfulness; irritability; appetite changes; sleep disturbances; social avoidance; alcohol and substance misuse.	Stress is a natural response to harmful situations (real or perceived) 'Fight-or-flight' response causes increased heart rate, breathing rate, blood pressure and mobilisation of substrate stores i.e. the release of stored energy (glucose, fats, and sometimes proteins) into the bloodstream so the body has enough fuel to respond to stress. Chronic stress can contribute to disease development.	Potential causes: Stress is multifactorial and may arise from work pressures, major life events, relationship difficulties, financial concerns, social isolation and health problems. Modifiable risk factors: poor diet, lack of exercise, poor sleep, ineffective coping strategies, alcohol or substance misuse, limited social support and lack of stress-management techniques or therapy. Non-modifiable risk factors: chronic illness, previous trauma or adverse childhood experiences, genetic predisposition, age and certain socioeconomic circumstances or caring responsibilities.

General Anxiety Disorder (GAD)

Signs and symptoms	Pathophysiology	Risks and causes
At least 3 of the following symptoms: Restlessness; nervousness; fatigue; poor concentration; irritability; muscle tension; sleep disturbances	Excessive worry that is disproportionate to any inherent risk present, occurring most days for at least six months	If untreated or unmanaged risk of: CVD (lifestyle factors) anxiety disorders, eating disorders; depression, substance misuse; self-harm risk; suicide Potential causes: childhood adversity or trauma, bullying or abuse, bereavement, relationship difficulties, unemployment and major life changes or transitions. Modifiable risk factors: chronic stress, poor coping strategies, substance misuse, poor sleep, social isolation and limited social support. Non-modifiable risk factors: family history/genetic predisposition, previous trauma or adverse childhood experiences, chronic physical illness and increasing age.

Appendix B: Medical interventions, medication side-effects and lifestyle interventions

Hypertension

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Antihypertensive e.g. ACE (angiotensin-converting enzyme) inhibitors	Reduce blood pressure by relaxing blood vessels	Dry tickly cough; dizziness; headaches; diarrhoea; itching or mild rash; blurred vision	• Increase physical activity and exercise • Reduce salt intake • Reduce alcohol intake • Smoking cessation • Relaxation (Benson method)
Calcium channel blockers	Dilate blood vessels to lower blood pressure	Headaches; swollen ankles; constipation	
Angiotensin-2 receptor blockers (ARBs)	Alternative to ACE inhibitors; reduce blood pressure by relaxing blood vessels	Dizziness; headaches; cold or flu-like symptoms	
Diuretics	Flush excess water and salt from the body through urine	Dizziness/light-headedness when standing; increased thirst; increased frequency of urination; rash	
Beta-blockers	Reduce blood pressure by making the heart beat slower and with less force	Fatigue; dizziness/light-headedness (can be a sign of low heart rate); cold fingers or toes (beta blockers can affect blood supply to hands and feet); difficulty sleeping/nightmares; nausea	

Hypercholesterolemia

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Statins	Reduce LDL cholesterol levels by reducing LDL produced by the liver	Indigestion; headaches; nausea; muscle aches	• Increase physical activity and exercise • Dietary changes – BHF guidance, NHS choices guidance • Smoking cessation

Hyperlipidaemia

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Statins	Reduce LDL cholesterol levels by reducing LDL produced by the liver	Indigestion; headaches; nausea; muscle aches	• Increase physical activity and exercise • Dietary changes – BHF guidance, NHS choices guidance • Smoking cessation
Cholesterol absorption inhibitors	Block dietary cholesterol being absorbed into the body	Diarrhoea; GI discomfort; GI disturbances; headache	
PCSK9 inhibitors	Lower cholesterol by blocking the receptors on the protein (PSKS9) in the liver which increases cholesterol levels, allowing the liver to clear away more cholesterol	Fatigue; muscle pain/soreness; swelling near injection site	
Bempedoic acid	Similar effect to statins	Anaemia; gout; pain in extremities	

Coronary Heart Disease

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Blood thinning medication (Aspirin)	Reduces the risk of a heart attack by thinning blood and preventing clotting	Mild indigestion; bleeding more easily than usual	• Increase physical activity and exercise • Dietary changes – BHF guidance, NHS choices guidance • Smoking cessation
Blood thinning medication (Clopidogrel)		Bleeding more easily than usual; diarrhoea; stomach pain; indigestion; heartburn	
Blood thinning medication (Rivaroxaban)		Bleeding more easily than usual; fatigue; shortness of breath	
Statins	Lower cholesterol	Indigestion; headaches; nausea; muscle aches	

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Beta-blockers	Lower blood pressure	Fatigue; dizziness/light-headedness (can be a sign of low heart rate); cold fingers or toes (beta blockers can affect blood supply to hands and feet); difficulty sleeping/nightmares; nausea	
Nitrates	Dilate blood vessels, allowing more blood to move through. Lower blood pressure and relieve heart pain	Headaches; dizziness; flushed skin	
Antihypertensive e.g. ACE (angiotensin-converting enzyme) inhibitors	Lower blood pressure	Dry tickly cough; dizziness; headaches; diarrhoea; itching or mild rash; blurred vision	
Angiotensin-2 receptor blockers (ARBs)	Lower blood pressure	Dizziness; headaches; cold/flu-like symptoms	
Calcium channel blockers	Lower blood pressure	Headaches; swollen ankles; constipation	

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Diuretics (water pills)	Remove excess water and salt	Dizziness/light-headedness when standing; increased thirst; increased frequency of urination; rash	

Peripheral Vascular Disease

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Blood thinning medication (Aspirin)	Reduces the risk of a heart attack by thinning blood and preventing clotting	Mild indigestion; bleeding more easily than usual	• Supervised exercise – increase physical activity and exercise • Dietary changes – BHF guidance, NHS choices guidance • Smoking cessation • Weight management • Reducing alcohol intake
Blood thinning medication (Clopidogrel)		Bleeding more easily than usual; diarrhoea; stomach pain; indigestion; heartburn	
Blood thinning medication (Rivaroxaban)		Bleeding more easily than usual; fatigue; shortness of breath	

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Statins	Lower cholesterol	Indigestion; headaches; nausea; muscle aches	
Antihypertensive e.g. ACE (angiotensin-converting enzyme) inhibitors	Lower blood pressure	Dry tickly cough; dizziness; headaches; diarrhoea; itching or mild rash; blurred vision	

Obesity

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Orlistat	Aids weight loss by reducing fat absorption	GI symptoms (abdominal discomfort/pain; faecal urgency; faecal incontinence; fatty stools); headache; respiratory infections; UTIs; hypoglycaemia; anxiety; fatigue; gingival; menstrual disturbances	• Increase physical activity and exercise • 30 minutes physical activity daily • Reduce sedentary time • Dietary changes - Increase fruit and vegetables • Smoking cessation • Counselling/talking therapies

Type 1 Diabetes

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Long-acting insulin (e.g. Detemir)	Taken 1-2 times per day. Increase insulin levels.	Hypoglycaemia; injection-site irritation; bruising; fatty lumps; visual changes	• Increase physical activity and exercise to meet CMO guidelines ○ Develop and maintain strength in major muscle groups • Dietary changes – Diabetes UK guidance, NHS choices guidance ○ Healthy balanced diet low fat, sugar and salt, at least 5 portions of fruit and veg a day ○ Low-GI diet is not recommended for type 1 diabetes – carbohydrate counting recommended (NB: dietary advice is outside scope of practice – client will need to work with a dietician) • Smoking cessation • Weight management • Reducing alcohol intake
Rapid/short-acting insulin	Taken before meals. Increases insulin levels		

Type 2 Diabetes

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Metformin	Lowers blood glucose by improving insulin sensitivity	Nausea; vomiting; diarrhoea; stomach ache; appetite loss; metallic taste in mouth.	• Dietary changes - Diabetes UK guidance, NHS choices guidance ○ High-fibre, low-GI carbohydrate sources ○ Healthy balanced diet - low in trans-fatty acids, sugar and salt • Smoking cessation • Reduce alcohol intake • Weight management • Weight loss if required • Minimise sedentary behaviour • Increase physical activity and exercise • Monitor blood glucose levels before exercise, follow recommendations for levels.

Hyperthyroidism

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Beta-blockers (e.g. propranolol, metoprolol, nadolol)	Control palpitations, tremor, tachycardia and anxiety	Fatigue; dizziness/light-headedness (can be a sign of low heart rate); cold fingers or toes (beta blockers can affect blood supply to hands and feet); difficulty sleeping/nightmares; nausea	• Smoking cessation, • Limiting caffeine intake if symptomatic • maintaining adequate nutrition and hydration • attending regular monitoring appointments • Adhering to prescribed treatment.
Carbimazole	Reduce thyroid hormone production	Nausea; vomiting; dizziness; headache; joint pain; itchy skin or rash	

Hypothyroidism

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Levothyroxine	Raise thyroxine levels	Nausea; vomiting; diarrhoea; headaches; restlessness; insomnia; flushing; sweating; muscle cramps; shaking	• Take levothyroxine consistently as prescribed, (ideally on an empty stomach and at the same time each day) • Avoid taking iron or calcium supplements close to the dose • Attend regular thyroid function monitoring • Maintain a healthy diet and regular physical activity.

Asthma

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Salbutamol (SABA)	Relaxes airway muscles	Feeling shaky; increased heart rate; headaches	• Exercise • Minimise exposure to triggers • Smoking cessation
Inhaled corticosteroids (ICS)	Reduce inflammation	Sore throat; hoarse or croaky voice; cough; oral thrush	

Chronic Obstructive Pulmonary Disease (COPD)

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Beta 2 agonist – Short-acting bronchodilator (SABAs) e.g. Salbutamol	Relaxes airway muscles	Feeling shaky; increased heart rate; headaches	• Exercise • Long-term oxygen therapy • Smoking cessation
Beta 2 agonist – Long-acting bronchodilator (LABAs) e.g. salmeterol, formoterol and indacaterol	Relaxes airway muscles	Arrhythmias; headaches; palpitations; tremor; muscle cramps	
Short-acting muscarinic antagonist (SAMA) e.g. Ipratropium	Relaxes airway muscles	Dry mouth; nasal or throat irritation; headache, and nausea	
Long-acting muscarinic antagonist (LAMAs) e.g. Tiotropium	Relaxes airway muscles	Dry mouth; sore throat; cough; headaches	

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Inhaled corticosteroids (ICS) e.g. Formoterol with budesonide	Reduce inflammation	Sore throat; hoarse or croaky voice; cough; oral thrush	
Theophylline tablets	Relaxes smooth muscle and reduces sensitivity to allergens	Nausea; vomiting; headaches; insomnia; palpitations	

Lower back pain

Some pain relievers are addictive.

v	Desired effects	Side effects	Lifestyle interventions
Nonsteroidal Anti-Inflammatory Drugs (NSAIDs)	Reduce pain and inflammation	Gastrointestinal (GI) symptoms (nausea, abdominal pain, diarrhoea); asthma exacerbation	• Exercise programmes • Postural improvement • Functional movement retraining • Manual therapy • Physiotherapy • Ergonomic modifications e.g. chair height and sleep positions etc.
Codeine	Pain relief	Constipation; nausea; drowsiness; confusion; dizziness; dry mouth; headaches	
Paracetamol	Pain relief	Side effects from paracetamol are rare, but can include allergic reactions (such as a rash or swelling), flushing, low blood pressure, or a fast heartbeat	

Osteoarthritis

Some pain relievers are addictive.

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Topical NSAIDs	Pain relief	Skin rash; photosensitivity; asthma symptoms	• Regular exercise • Weight management • Weight loss if required • Suitable footwear • Physiotherapy
Corticosteroids	Reduce inflammation	Skin thickening; increased osteoporosis risk	
Paracetamol	Pain relief	Side effects from paracetamol are rare, but can include allergic reactions (such as a rash or swelling), flushing, low blood pressure, or a fast heartbeat	

Rheumatoid arthritis

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
NSAIDs	Pain relief	Gastrointestinal (GI) symptoms (nausea, abdominal pain, diarrhoea); asthma symptoms	• Activity during remission • Weight management • Weight loss if required • Physiotherapy
Conventional disease modifying anti-rheumatic drug (cDMARDs) e.g. methotrexate, leflunomide, sulfasalazine, hydroxychloroquine	Suppress immune attack on joints	Gastrointestinal (GI) symptoms (nausea, abdominal pain, diarrhoea)	
Glucocorticoids	Anti-inflammatory and immunosuppressive	Weight gain; fluid retention; increased hunger; mood swings; blurred vision; insomnia; muscle weakness; feeling nervous; restlessness	

Osteopenia

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Calcium and Vitamin D supplements	Provide daily calcium and vitamin D Support bone health	Constipation; flatulence; abdominal pain; diarrhoea; hypercalciuria; skin rash	• Regular exercise • Increase calcium intake • Increase vitamin D intake • Reduce caffeine • Reduce alcohol • Stop smoking

Osteoporosis

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Calcium and cholecalciferol (vitamin D3)	Provide daily calcium (1g) and vitamin D (400 units) Support bone health	Constipation; flatulence; abdominal pain; diarrhoea; hypercalciuria; skin rash	• Regular exercise • Weight-bearing exercise • Increase calcium intake • Increase vitamin D intake • Reduce caffeine intake • Reduce alcohol intake • Smoking cessation
Bisphosphonates	Increase bone mineral density by altering osteoclast activity and function, inhibiting bone reabsorption	Nausea; dyspepsia (chronic indigestion); gastritis; abdominal pain; bone pain; joint pain; muscle pain	
Hormone Replacement Therapy (HRT)	Maintain bone density in postmenopausal women	Breast tenderness, bloating, nausea, headaches, and breakthrough bleeding	

Joint replacement

Some pain relievers are addictive.

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Nonsteroidal Anti-Inflammatory Drugs (NSAIDs)	Reduce pain and inflammation	Gastrointestinal (GI) symptoms (nausea, abdominal pain, diarrhoea); asthma exacerbation	• Exercise to strengthen muscles around the joint • Improve mobility
Codeine	Pain relief	Constipation; nausea; drowsiness; confusion; dizziness; dry mouth; headaches	
Paracetamol	Pain relief	Side effects from paracetamol are rare, but can include allergic reactions (such as a rash or swelling), flushing, low blood pressure, or a fast heartbeat	

Depression

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Antidepressants: Selective serotonin inhibitor (SSRIs)	SSRIs increase serotonin levels in the brain by preventing it from being reabsorbed too quickly. This allows serotonin to remain available for longer, helping to improve mood and emotional regulation.	Agitation; anxiety; diarrhoea; nausea and vomiting; dizziness; blurred vision; reduced libido	• Increase physical activity • Dietary changes • Reduce caffeine • Reduce alcohol • Relaxation and mindfulness • Cognitive Behavioural Therapy (CBT)
Antidepressants: Tricyclic	Increase serotonin and noradrenaline in the brain chemicals which regulate mood, circadian rhythms and memory	Dry mouth; blurred vision; constipation; urinary issues; sweating; light-headedness; excessive drowsiness	
Antidepressants: Serotonin-noradrenaline reuptake inhibitor (SNRIs)	Increase serotonin and noradrenaline	Nausea; headaches; drowsiness; dizziness; dry mouth; constipation; insomnia; reduced libido; increased blood pressure	

Stress

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Lifestyle and non-pharmaceutical interventions	UK doctors usually recommend lifestyle changes and talking therapies (such as Cognitive Behaviour Therapy) first to reduce stress, keeping medication as an option to manage symptoms if these do not help	N/A	• Dietary changes –NHS choices guidance • Reduce caffeine intake • Reduce alcohol intake • Cognitive Behavioural Therapy (CBT) • Relaxation and mindfulness

General Anxiety Disorder (GAD)

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Antidepressants: Selective serotonin inhibitor (SSRIs)	SSRIs increase serotonin levels in the brain by preventing it from being reabsorbed too quickly. This allows serotonin to remain available for longer, helping to improve mood and emotional regulation.	Agitation; anxiety; diarrhoea; nausea and vomiting; dizziness; blurred vision; reduced libido	• Increase physical activity • Dietary changes • Reduce caffeine • Reduce alcohol

Medical/Intervention	Desired effects	Side effects	Lifestyle interventions
Antidepressants: Serotonin-noradrenaline reuptake inhibitor (SNRIs)	Increase serotonin and noradrenaline	Nausea; headaches; drowsiness; dizziness; dry mouth; constipation; insomnia; reduced libido; increased blood pressure	• Relaxation and mindfulness • Cognitive Behavioural Therapy (CBT)
Pregabalin	Reduces release of anxiety-inducing brain chemicals	Headaches; drowsiness; mood changes; nausea; swelling of extremities; blurred vision; weight gain (due to increased hunger); memory problems	

Appendix C: Exercise guidelines and information sources

Hypertension

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Normalise blood pressure; reduce CHD risk	Low-to-moderate intensity exercise; extended warm-up and cool-down; avoid Valsalva manoeuvre; use RPE if medication affects heart rate response; slow transitions from floor to standing; promote correct breathing; include relaxation; consider medication side effects and other CVD risk factors	British Heart Foundation: https://www.bhf.org.uk British and Irish Hypertension society: https://bihsoc.org

Hyperlipidaemia

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Improve blood lipid profile; improve HDL:LDL ratio; reduce CHD risk	Consider other CVD risk factors and comorbidities, medication side effects and other cardiovascular risk factors	British Heart Foundation: https://www.bhf.org.uk Heart UK: https://www.heartuk.org.uk

Hypercholesterolaemia

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Improve cholesterol management; increase HDL:LDL ratio; reduce CHD risk	Consider CVD risk factors and comorbidities; consider medication side effects low intensity and resistance initially; extended warm-up and cool-down	British Heart Foundation: https://www.bhf.org.uk Heart UK: https://www.heartuk.org.uk

Coronary Heart Disease

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Improve circulatory efficiency; lower cholesterol; normalise blood pressure; support weight management; reduce heart attack risk	Consider other CVD risk factors; monitor exercise intensity closely (work within medically prescribed restrictions); start at low intensity (40–60% HRR) and increase gradually over time; low-load, higher-repetition resistance training; clients with angina should carry glyceryl trinitrate spray (GTN spray) or nitroglycerin tablets;	British Heart Foundation: https://www.bhf.org.uk British Association for Cardiovascular prevention and Management (BACPR): https://www.bacpr.org/

Peripheral Vascular Disease

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Improve circulation; lower cholesterol; normalise blood pressure; aid weight management	Ensure comorbidities are stable, well controlled and within scop; low-intensity aerobic activity; upright activities such as walking (to encourage blood flow to lower body); interval approach if claudication (pain/cramping/aching in limbs) occurs; low-load, higher-repetition upper body resistance training; suitable footwear and proper foot care; consider medication side effects	The Circulation Foundation: https://www.circulationfoundation.org.uk

Obesity

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Aids weight management; supports weight loss; reduces risk of CHD, stroke and Type 2 diabetes; strengthen joints; improves mental and emotional wellbeing	Consider stress on joints and body systems; functional limitations; bulk and size, emotional sensitivities; comorbidities (CHD, diabetes); low-impact and low-weight-bearing activities; promote activities of daily living (ADL)	Obesity UK: https://www.obesityuk.org.uk

Type 1 Diabetes

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Aids weight management; supports weight loss; reduces blood pressure; reduces risk of CHD	Must be stable and controlled; must be able to self-monitor blood glucose throughout activity; avoid exercise if BG >13 mmol/L or <4 mmol/L until restored; do not exercise if hypo or severe hypo in last 24 hours; carry hypo treatment and carb snack(s); may need to adjust medication dosage prior to exercise; monitor for hyper/hypoglycaemia; consider side effects of medication; consider insulin timing and injection sites relative to body part taking part in exercise; ensure suitable footwear	Diabetes UK: https://www.diabetes.org.uk DAFNE: https://dafne.nhs.uk

Type 2 Diabetes

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Improve insulin sensitivity; weight management; reduce blood pressure; reduce CHD risk	Similar considerations to Type 1 diabetes, including monitoring blood glucose before and after exercise and recognising symptoms of hypoglycaemia; however, exercise is particularly important in managing insulin resistance and supporting weight management in Type 2 diabetes.	Diabetes UK: https://www.diabetes.org.uk

Hyperthyroidism

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Improve mood; improve sleep; improve bone density; increase energy; reduce CHD risk; relieves other symptoms of thyroid disease.	Condition must be controlled; begin with low-to-moderate intensity exercise; consider elevated metabolism and heart rate of hyperthyroidism (which if uncontrolled can put excess strain on heart) and medication effects	British Thyroid Foundation: https://www.btf-thyroid.org

Hypothyroidism

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Improve mood; improve sleep; improve bone density; increase energy; reduce CHD risk; boosts metabolism; aids weight management; supports weight loss; relieves other symptoms of thyroid disease	Condition must be controlled; begin at low-to-moderate intensity exercise consider slower metabolism and heart rate.; weight loss may progress more slowly (set realistic goals); consider medication effects	British Thyroid Foundation: https://www.btf-thyroid.org

Asthma

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Strengthen respiratory muscles; improve cardiovascular fitness and function; improve expiration; reduces risk of CHD.	Asthma must be controlled; be aware of triggers and allergies; be aware of how to manage an asthma attack; regular GP checks required to monitor condition; prolonged warm-up and cool-down; interval approach preferred; monitor intensity closely – use breathlessness scale; warm humid environments may be beneficial (e.g. swimming); consider medication side effects	Asthma and Lung UK: https://www.asthmaandlung.org.uk

Chronic Obstructive Pulmonary Disease (COPD)

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Improve oxygen uptake; improve ADL; maintain function and independence	Consider respiratory muscles may be weakened; breathlessness; low exercise tolerance; limited ventilatory capacity; skeletal muscle disfunction; comorbidities; consider medication side effects and psychological factors; avoid temperature extremes; emphasise posture training; breathing exercises; strengthen lower limbs; include trunk and spinal mobility; promote relaxation.	Asthma and Lung UK: https://www.asthmaandlung.org.uk

Lower back pain

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Improve posture; improve functional movement and ability; strengthen core; maintain mobility and flexibility; maintain functional movement, reduce risk of CHD	Avoid heavy lifting, jarring, repetitive bending and twisting; avoid high impact; promote posture and abdominal engagement; avoid prolonged static positions; promote correct lifting; during episodes of LBP focus on abdominal hollowing rather than flexion exercises; consider emotional impact and medication side effects	Back care: https://backcare.org.uk Versus arthritis: https://www.versusarthritis.org – Leaflet on back pain exercises – Back pain exercise sheet Chartered Society of Physiotherapists: https://www.csp.org.uk

Osteoarthritis

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Boost synovial fluid; maintain range of movement (ROM); stabilise affected joints; aids weight loss; maintain functional movement	Consider pain and discomfort particularly with certain exercises (e.g. kneeling, weight bearing on wrists); avoid prolonged exercise in one position; avoid excessive impact; promote mobility and flexibility; strengthen around affected joints; use suitable footwear; consider side effects of medication	Versus arthritis: https://www.versusarthritis.org Arthritis action: https://www.arthritisaction.org.uk

Rheumatoid arthritis

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Maintain ROM; stabilise affected joints during remission; aids weight loss; maintain functional movement; reduce risk of CHD	No exercise during flare-ups; consideration to previous joint damage; avoid excessive impact, repetition and contact sports; avoid fast paced movements; morning stiffness may affect exercise timing; consider emotional effects and medication side effects	National Rheumatoid Arthritis Society: https://nras.org.uk Versus arthritis: https://www.versusarthritis.org

Osteopenia

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Maintain bone density; reduce falls risk; maintain functional movement; reduce risk of CHD	Promote weight-bearing activity; bone loading; strengthening exercises; strengthen core to support spine; promote ADLs; consider medication side effects	Royal Osteoporosis Society: https://theros.org.uk

Osteoporosis

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Maintain bone density; reduce falls risk; maintain functional movement; reduce risk of CHD	Disease severity must be considered; environmental safety is essential; clients may fear falling; promote weight-bearing activity; bone loading; strengthening exercises; promote ADLs; consider medication side effects	Royal Osteoporosis Society: https://theros.org.uk

Joint replacement

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Improve functional movement; improve mobility; strengthen muscles surrounding the new joint; reduce risk of CHD	Hip: observe surgeon-specific precautions regarding hip flexion, adduction and rotation during the postoperative period; monitor for signs of dislocation; ensure seated exercise positions keep hips above knee height where advised. Knee: initially favour low-impact activities and avoid high-impact exercise or twisting movements until cleared by the orthopaedic and physiotherapy team; progression should be individualised according to recovery and prosthesis type	NHS https://www.nhs.uk/tests-and-treatments/hip-replacement/recovering-from-a-hip-replacement/ https://roh.nhs.uk/news-events/enhancing-recovery-following-total-knee-replacement

Depression

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Improves mood; self-esteem; self-worth; confidence; body image; cognitive function; maintain functional movement reduces anxiety and risk of CHD	Low energy and low motivation common; lethargy; encourage outdoor exercise in sunshine if possible; quieter environments/quieter times of day may improve adherence and enjoyment when client is more comfortable; promote ADLs; consider medication side effects and comorbidities	MIND: https://www.mind.org.uk Mental Health Foundation: https://www.mentalhealth.org.uk Royal College of Psychiatrists: https://www.rcpsych.ac.uk – Patient information leaflets

Stress

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Improves mood; self-esteem; self-worth; confidence; body image; cognitive function; maintain functional movement reduces anxiety and risk of CHD	Promote relaxation techniques and breathing exercises; consider medication and co-morbidities	MIND: https://www.mind.org.uk Mental Health Foundation: https://www.mentalhealth.org.uk Royal College of Psychiatrists: https://www.rcpsych.ac.uk – Patient information leaflets

General Anxiety Disorder

Benefits of exercise	Exercise considerations Risks and restrictions	Sources
Improves mood; self-esteem; self-worth; confidence; body image; cognitive function; maintain functional movement reduces anxiety and risk of CHD	Explain that elevated heart rate, sweating and breathlessness are normal exercise responses; gradual warm-up; quieter exercise environments may help; promote relaxation and breathing; consider medication side effects and co-morbidities	MIND: https://www.mind.org.uk Mental Health Foundation: https://www.mentalhealth.org.uk Royal College of Psychiatrists: https://www.rcpsych.ac.uk – Patient information leaflets

Additional sources of information


NICE	https://www.nice.org.uk
NHS	https://www.nhs.uk
Patient UK	https://patient.info
SIGN	https://www.sign.ac.uk (part of Healthcare Improvement Scotland)
HealthTalk	https://healthtalk.org
ACSM	https://www.acsm.org
World Health Organization (WHO)	https://www.acsm.org

Guidance for training providers

Centre and qualification approval

Before you can begin delivery of this qualification, you must be a Y Awards Ltd centre with appropriate qualification and staff approval.

Find out more on our website:




ymcaawards.co.uk/approvals

Tutor, assessor, and IQA requirements

All tutors, assessors and internal quality assurance (IQA) staff need to hold:

- a subject matter qualification.
- a qualification related to the role that they will be performing (tutor, assessor, or IQA).

Find out more on our website:



ymcaawards.co.uk/approvals/staff-approval

Blank Page



Y Awards Ltd, has a long and respected history as the first UK awarding organisation to specialise in health and fitness qualifications, launching in 1998 as CYQ.

We're experts in education, health, and wellbeing with over 20 years of experience developing UK-regulated and globally-recognised qualifications. We work closely with industry experts, employers and training providers to make sure that our products and services deliver life-changing opportunities.

With over half a million qualifications awarded, over 400,000 people have advanced their careers with Y Awards Ltd.



www.ymcaawards.co.uk